

Modulation of dendritic cell function by the fungal quorum sensing molecule farnesol

Dissertation

To fulfill the
requirements for the degree of
„doctor rerum naturalium“ (Dr. rer. nat.)

**Submitted to the Council of the Faculty
of Biological Sciences
of the Friedrich Schiller University Jena**

**by Wolfgang Vivas
born on January 18, 1990
in Caracas, Venezuela**



FRIEDRICH-SCHILLER-
UNIVERSITÄT
JENA

The work included in this thesis was carried out in the Septomics Research Center from the Friedrich Schiller University Jena and Leibniz Institute for Natural Product Research and Infection Biology, Hans Knöll Institute-Jena, in the research group “Fungal Septomics” under the supervision of Prof. Dr. Oliver Kurzai



The PhD project was supported by a fellowship from the Jena School for Microbial Communication (JSMC)

Dissertation examiners

1. Prof. Dr. Oliver Kurzai, Julius-Maximilians University, Würzburg and Leibniz Institute for Natural Product Research and Infection Biology - Hans-Knöll Institute, Jena, Germany.
2. Prof. Dr. Marc Thilo Figge, Leibniz Institute for Natural Product Research and Infection Biology - Hans-Knöll Institute, Jena, Germany.
3. Prof. Dr. Doris Wilflingseder, Medical University of Innsbruck, Austria.

Date of public defense: 23.06.2020

Table of Contents

1	Summary.....	1
2	Zusammenfassung.....	2
3	Introduction	3
3.1	<i>Candida albicans</i>	3
3.2	Farnesol production by <i>Candida albicans</i>	4
3.2.1	Effects of farnesol on <i>Candida albicans</i>	4
3.2.2	Farnesol effects on other microorganisms	5
3.2.3	Farnesol effects on mammalian cells	6
3.3	Role of dendritic cells in anti- <i>Candida</i> immune responses	7
3.3.1	Dendritic cells	7
3.3.2	DC antigen presentation and T cell polarization	10
3.3.3	DC signaling and immunity against <i>Candida albicans</i>	11
3.3.4	Nuclear receptors and their role in DC maturation and immune response against fungal infections	13
3.3.4.1	Nuclear receptors	13
3.3.4.1.1	Retinoid X receptor: RXR	15
3.3.4.1.2	Retinoic acid receptor: RAR	16
3.3.4.1.3	Peroxisome proliferator activated receptors: PPAR	17
3.3.4.1.4	Liver X receptor: LXR	18
4	Aims of the work	21
5	Experimental procedures	22
5.1	Reagent preparation	22
5.2	Dendritic cell generation and maturation	22
5.3	<i>Candida albicans</i> cultivation	23
5.4	Flow cytometry	24
5.4.1	Viability determination	24
5.4.2	Characterization of cell surface markers	24
5.5	Invariant Natural Killer T cells expansion	25
5.6	Intracellular staining	25
5.7	Phosphoflow	26
5.8	Cytokine measurement	26

5.9	RNA isolation and quantitative real-time PCR	26
5.10	Statistical analysis	27
6	Results	28
6.1	Viability of monocyte-differentiated DC in presence of farnesol	28
6.2	Farnesol-differentiated DC have altered activation in response to distinct stimulation	29
6.3	Molecular mechanisms modulated by farnesol on DC differentiation and maturation.....	33
6.3.1	Differentiation of DC in presence of farnesol increased the expression of genes involved in nuclear receptor activity	33
6.3.1.1	Activation of PPAR γ and RAR α leads to a DC phenotype comparable to farnesol-differentiated DC	36
6.3.1.2	Nuclear receptor inhibitors have no significant effect on dendritic cell viability	38
6.3.1.3	Farnesol modulates DC phenotype partially through activation of the PPAR γ and RAR α signaling pathways	39
6.3.1.4	Release of cytokines and chemokines by farnesol-differentiated DC is mainly independent of nuclear receptors activation	45
6.3.2	Role of MAPK and NF- κ B signaling cascades in farnesol-differentiated DC	47
6.3.2.1	MAPK and NF- κ B signaling pathways modulate maturation of DC	47
6.3.2.2	Farnesol regulates cytokine and chemokine secretion through MAPK and NF- κ B signaling pathways	49
6.4	Functional impact of farnesol treatment in DC	53
6.4.1	Farnesol-differentiated DC have a lower capacity to activate iNKT and Th1 cells ...	53
6.4.1.1	Altered secretion of IL-12 and IL-10 influences the capacity of farnesol-differentiated DC to activate different T cell subsets.....	56
6.4.2	Farnesol-differentiated DC have a higher expression of inhibitory receptors but lack capacity to induce regulatory T cells expansion	59
7	Discussion	62
7.1	Farnesol impairs DC differentiation and maturation partially through nuclear receptors activation	62
7.2	Farnesol regulation of CD1 molecules and its functional impact	63
7.3	Farnesol alters cytokines release through several pathways	65
7.4	Farnesol-treated DC have a lower capacity to activate different T cell subsets	66
8	Conclusions.....	68
9	References.....	70

10	List of abbreviations.....	83
11	Index of tables and figures	85
12	Appendix	87
12.1	Curriculum Vitae.....	87
12.2	Acknowledgments	90
12.3	Declaration of originality	92

1 Summary

Microorganisms communicate with each other through the secretion of molecules that accumulate in the environment and mediate specific responses upon reaching a specific threshold. This process has been termed quorum sensing and is essential to coordinate changes in behavior dependent on cell population density. Farnesol (FOH) is the first quorum sensing molecule described in a eukaryotic organism, and it is produced by the opportunistic pathogenic fungus *Candida albicans*. FOH regulates morphogenesis in *C. albicans*; however, we and others have described a new role of FOH as a compound with immunomodulatory properties. In particular, FOH showed to impair DC differentiation from monocytes, resulting in altered expression of surface markers and secretion of cytokines, thereby suppressing DC ability to induce T cell proliferation. However, the molecular mechanisms by which FOH modulates DC differentiation and maturation are poorly understood. The results of this study revealed that FOH regulates DC function through several signaling pathways. FOH modulates the expression of costimulatory molecules partially through activation of the nuclear receptors PPAR γ , RAR α and LXR α . In particular, FOH increased the expression of the lipid antigen-presenting molecule CD1d through activation of p38 MAPK and PPAR γ /RAR α signaling pathway. However, CD1d upregulation did not confer these cells an elevated capacity to activate invariant Natural Killer T (iNKT) cells. FOH-differentiated DC showed diminished secretion of IL-12 and increased IL-10 release. Interestingly, reconstitution of the IL-12/IL-10 cytokine milieu restored FOH-differentiated DC to activate iNKT, Th1 and regulatory T cells. Mechanistically, FOH modulates IL-10 and pro-inflammatory cytokines through activation of RAR α , LXR α , MAPK and NF- κ B signaling pathways. Altogether, our results showed that FOH induces paralysis of DC function through activation of nuclear receptors, MAPK and NF- κ B signaling pathways. Since these cells play an important role in regulating the immune response during infection, this work supports the role of FOH as a virulence factor produced by *C. albicans* to overcome immune surveillance by DC.

2 Zusammenfassung

Mikroorganismen kommunizieren miteinander durch die Sekretion von Molekülen, die sich in der Umwelt ansammeln und bei Erreichen eines bestimmten Schwellwertes bestimmte Reaktionen vermitteln. Dieser Prozess wird als Quorum Sensing bezeichnet und dient dazu, Verhaltensänderungen in Abhängigkeit der Zellpopulationsdichte zu koordinieren. Farnesol (FOH) ist das erste Quorum-Sensing-Molekül, das in einem eukaryotischen Organismus beschrieben wurde. Es wird von dem opportunistischen und pathogenen Pilz *Candida albicans* produziert und dient der Regulation der Morphogenese. Wir und andere haben eine neue Funktion von FOH beschrieben, in der es als Verbindung mit immunmodulatorischen Eigenschaften dient. FOH beeinträchtigte die Differenzierung von Monozyten zu Dendritischen Zellen (DZ), was zu einer veränderten Expression von Oberflächenmarkern und zur Sekretion von Zytokinen führte, wodurch die Fähigkeit der DZ zur Induktion der T-Zell-Proliferation unterdrückt wurde. Die molekularen Mechanismen, über die FOH die Differenzierung und Reifung von DZ moduliert sind jedoch kaum bekannt. Die Ergebnisse dieser Studie zeigten, dass FOH die Funktion von DZ über mehrere Signalwege reguliert. FOH moduliert die Expression kostimulatorischer Moleküle teilweise durch Aktivierung der Nuklearrezeptoren PPAR γ , RAR α und LXR α . FOH erhöhte insbesondere die Expression des Lipidantigen-präsentierenden Moleküls CD1d durch Aktivierung des p38-MAPK- und PPAR γ /RAR α -Signalweges. Die Hochregulation von CD1d verlieh diesen Zellen jedoch keine erhöhte Fähigkeit, invariante natürliche Killer T (iNKT)- Zellen zu aktivieren. FOH-differenzierte DZ zeigten eine verminderte IL-12 und eine erhöhte IL-10 Sekretion. Interessanterweise befähigte die Rekonstitution des IL-12/IL-10-Zytokinmilieus die FOH-differenzierten DZ wieder, iNKT-, Th1- und regulatorische T-Zellen zu aktivieren. Mechanistisch modulierte FOH die Freisetzung von IL-10 und entzündungsfördernden Zytokinen durch Aktivierung von RAR α -, LXR α -, MAPK- und NF- κ B-Signalwegen. Insgesamt zeigten unsere Ergebnisse, dass FOH die Paralyse der DZ durch Aktivierung der Nuklearrezeptoren und über MAPK- und NF- κ B -Signalwege induzierte. Da diese Zellen eine wichtige Rolle bei der Regulierung der Immunantwort bei Infektionen spielen, unterstützt diese Arbeit die Rolle von FOH als Virulenzfaktor, der von *C. albicans* produziert wird, um die Immunüberwachung durch DZ zu überwinden.

3 Introduction

3.1 *Candida albicans*

C. albicans is one of the most prominent fungal commensal and pathogen of humans. *C. albicans* colonizes the gastrointestinal tract, mouth, skin and the female reproductive tract (1, 2). To colonize these niches, the fungus is able to adapt to several environmental conditions, including oxygen levels, host microbiome or changes in pH (1). However, *C. albicans* can become pathogenic if the host develops specific risk factors, such as microbial dysbiosis, immunodeficiency, major surgery, the presence of venous catheter and antibiotics (2, 3). Indeed, *C. albicans* can cause a broad range of infections in human, including superficial infections, such as oral or vaginal candidiasis, and life-threatening systemic infections (1). *C. albicans* relies on several virulence factors, including the expression of adhesins and biofilm formation, secretion of cytolytic proteins and morphological plasticity (4-10). *C. albicans* is a polymorphic fungus that can grow as budding yeast, pseudophyphae and true-hyphae (1, 2). The yeast form is associated with colonization and dissemination, while the filamentous form is involved in adhesion, penetration and tissue invasion (5, 6). Different environmental conditions are known to influence morphological transition in *C. albicans*, such as pH, starvation, the presence of serum or N-acetylglucosamine, temperature and CO₂ levels (6). Moreover, morphogenesis is also modulated by quorum sensing (QS), a mechanism of microbial communication to coordinate population density-dependent changes in behavior. In QS, secreted molecules from growing microorganisms accumulate in the local environment and modulate specific responses once a critical threshold or concentration has been reached. The main QS molecules produced by *C. albicans* include tyrosol, farnesoic acid and farnesol (1, 11-13). Tyrosol decreases the lag phase growth and stimulates filamentation as well as biofilm formation. However, its effects are only observed in absence of farnesol (11, 14). Farnesoic acid and farnesol inhibits the switch from yeast to hyphae form in a growing population. However, the effect of farnesoic acid is less than farnesol and only produced by one strain of *C. albicans* (12).

3.2 Farnesol production by *Candida albicans*

Farnesol (FOH) is a QS molecule produced mostly by *C. albicans* and inhibits the yeast to hyphae transition, which is one of the most important virulence factor associated with this fungus (1, 15). FOH is produced by *C. albicans* as a side product of the sterol biosynthetic pathway, which is essential for the membrane integrity of the fungus (16). Indeed, FOH is an acyclic sesquiterpene (isoprenoid/ 1-hydroxy-3, 7, 11-trimethyl-2, 6, 10-dodacatrien) produced by dephosphorylation of farnesol pyrophosphate (FPP), a key intermediate of the mevalonate pathway (17). Inhibition of the sterol synthetic pathway downstream of FOH induces accumulation of FPP, resulting in elevated concentration of FOH compared to untreated cells (18). Furthermore, inhibition of 3-hydroxy-methyl-glutaryl-CoA reductase (HMGR), necessary for mevalonate synthesis, reduces FOH production in *C. albicans* (19). Of the four existing isomers, only *E-E* FOH showed QS activity (20). Interestingly, FOH production is not unique to *C. albicans*. A study testing the capability of eight *Candida* species to produce FOH showed that *C. dubliniensis* can produce this QS molecule, but in much less quantities compared to *C. albicans* (21).

Production of FOH by *C. albicans* is likely dependent on the coordinated activation of several genes. Particularly, *DPP3* gene is known to be important for FOH biosynthesis in *C. albicans*. Production of FOH is significantly reduced in a *DPP3* deletion mutant (22). Furthermore, overexpression of *DPP3* increases the morphogenic-inhibitory effects of FOH (23). Furthermore, synthesis was enhanced in deletion mutants of the morphogenic repressor genes *NRG1* and *TUP1*, as well as in the associated hyphal-maintenance gene *EED1* (24-26).

3.2.1 Effects of farnesol on *Candida albicans*

The most prominent effect of FOH is its ability to influence *C. albicans* morphology, by blocking the transition from yeast to filamentous cell without altering the growth rate (27, 28). Additionally, FOH inhibits biofilm formation in *C. albicans*. Ramage *et al.* showed that the effect of FOH on biofilm formation depends on the concentration of the compound and the initial time of adherence (29). While low FOH concentration did not affect biofilm formation, incubation with 300 μ M FOH reduced *C. albicans* adherence and increased dispersal during the first 30 min of treatment. Interestingly, biofilm formation was impaired when cells were

incubated with 100 μ M FOH for 24 hours (30). These results indicate that FOH treatment might promote *C. albicans* dispersal to colonize new niches. In addition to morphogenesis and biofilm formation, FOH regulates the expression of genes involved in drug resistance, iron transport, cell wall maintenance, oxidative stress, heat shock, surface hydrophobicity and iron transport (30-32).

Several reports have shown that FOH impacts multiple signaling pathways in *C. albicans*. The Ras1-cAMP-PKA cascade is one of the first signaling pathways identified to be influenced by farnesol. Ras proteins are GTPases localized in the plasma membrane which plays an important role in filamentation (33). Piispanen *et al.* (34) showed that FOH cleave Ras1 from the plasma membrane and the resulting soluble Ras1 reduced the ability to activate the catalytic activity of the adenylyl cyclase Cyr1p, supporting the formation of yeast cells. One important event on Ras1 localization in the plasma membrane involves farnesylation of the protein, which facilitates Cyr1p binding to Ras1 (35, 36). Thus, FOH could also block Ras1 activity via inhibition of farnesylation (37). Interestingly, FOH alters several other signaling pathways, such as activation of the general stress mitogen-activated protein kinase (MAPK) Hog1 and the two components signaling via the Cek1-MAPK pathway and stabilization of transcription factors like Tup1/Nrg1, leading to repression of filamentation (15, 25, 38-40). Finally, FOH modulates the activation of Efg1 and Czf1 transcription factors activities resulting in the transition from white to opaque cells, with opaque cells being killed at lower FOH concentrations (15, 41).

3.2.2 Farnesol effects on other microorganisms

FOH does not only affect *C. albicans* itself but alters other microorganisms and cell types (15). FOH blocks conidiation and germination of *Aspergillus niger*, *Penicillium expansum*, and *Fusarium graminearum* via deregulation of G-protein/cAMP signaling pathway and induction of apoptosis. Interestingly, the induction of cell death by FOH in *Fusarium graminearum*, *Saccharomyces cerevisiae*, *A. nidulans* and *A. flavus* is a process dependent on metacaspase activation and ROS production (42-47). However, a different study found that FOH alters the membrane integrity of microorganisms due to its amphiphilic characteristic (48). Indeed, FOH treatment showed to blocks actin cytoskeleton formation in *A. fumigatus* by misplacing the hyphal tip localization of two prenylated Rho family GTPases,

AfRho1 and AfRho2, while inhibiting the cell wall integrity signaling pathway (48). Similarly, FOH increases drugs susceptibility of *Escherichia coli*, and *Staphylococcus aureus* via disruption of the cell membrane, resulting in the reduce formation of biofilms (49, 50).

3.2.3 Farnesol effects on mammalian cells

Besides FOH effects on *C. albicans* and other microorganisms, FOH has been shown to interact with mammalian cells. In particular, FOH induces apoptosis in several types of cancer cell lines, such as lung carcinoma cells, human oral squamous carcinoma cells, prostate cancer cells and breast cancer cells highlighting its relevance as a possible antineoplastic drug (51-54). However, some studies have found that FOH treatment also impacts viability of non-malignant cells. Particularly, FOH induces apoptosis and necrosis in spermatozoa (55). While FOH treatment reduced murine macrophages viability *in vitro*, the pro-apoptotic effect of this molecule on immune cells only occur at high concentrations (56, 57).

Although FOH inhibits filamentation in *C. albicans*, existing data suggest that FOH is likely a virulence factor promoting disease progression in systemic infections (22, 58). Treatment of *C. albicans* with subinhibitory concentrations of FOH-inducing azoles enhances virulence in a systemic candidiasis murine model (58). Furthermore, administration of FOH in mice infected with *C. albicans* reduced IL-12 and IFN- γ concentration, while increased kidney fungal burden and mortality (22). In accordance, infection with *C. albicans* strains that do not produce FOH showed reduced mortality in mice (22, 58). Likewise, treatment of mice with pravastatin, a HMG-CoA reductase inhibitor, decreased FOH production and enhanced viability during *C. albicans* infection (59). Interestingly, FOH treatment showed to be protective against oral candidiasis in mice.

Further evidences support immunomodulatory properties of FOH. Specifically, FOH treatment stimulates the chemokinesis of murine macrophages *in vitro* and *in vivo*, and these effects were enhanced in the presence of yeast cell wall components and aromatic alcohols (60). Interestingly, FOH administration with zymosan resulted in modulation of RAW264.7 macrophages genes expression, by enhancing IL-6, Toll-like receptor 2 (TLR2) and dectin-1 expression (61). However, Leonhardt *et al.* showed that FOH alters neutrophils, monocytes and dendritic cells functionality (56). FOH treatment promotes activation of

neutrophils and monocytes. However, this did not correlate with elevated phagocytic activity. In contrast, FOH dramatically impaired the differentiation of monocytes into dendritic cells. Particularly, FOH modified the expression of costimulatory and antigen-presenting molecules. These changes in surface molecules were accompanied by enhanced anti-inflammatory IL-10 and diminished IL-12 release and correlated with reduced capacity to induce T cell proliferation (56). Due to the increasing evidences of FOH as an important virulence factor and the relevance of dendritic cells in the anti-*Candida* immunity, this thesis focused in elucidating the molecular mechanisms in which FOH treatment orchestrates dendritic cells functionality.

3.3 Role of dendritic cells in anti-*Candida* immune responses

The innate immune system is the first-line of defense against invading pathogens, including *C. albicans*. While neutrophils and monocytes present one of the most important cellular components in the defense against this fungus, subsequent recognition and activation by dendritic cells leads to specific T cell anti-*Candida* immunity, which is important for fungal clearance (62, 63).

3.3.1 Dendritic cells

The orchestration of an effective immune response depends on dendritic cells (DC), a class of cells derived from bone-marrow that can be found in virtually most peripheral tissues, blood and lymphoid organs (64). DC are able to process extra- and intracellular antigens and present these via Major Histocompatibility Complex (MHC) molecules to T cells (65). Thus, DC are the bridge between the innate and adaptive immunity. DC population comprised of different types of cells, each with specific functions. They can be classified in plasmacytoid DC, myeloid conventional DC 1 (cDC1), myeloid conventional DC 2 (cDC2), Langerhans cells and monocyte-derived DC (64). Plasmacytoid DC are mainly present in blood and are specialized to sense and respond to viral infection by a rapid secretion of type I and type III interferons (66). Myeloid cDC1 are present in blood and tissues. These cells have the capacity to present antigens via MHC class I to promote CD8⁺ T cells and NK cells activation through IL-12 production (64, 67). Myeloid cDC2 are present in blood, tissues and lymphoid organs. These cells are equipped with a range of C-type lectins, Toll-like receptors

(TLR), Nucleotide-binding oligomerization domain (NOD)-like receptors and Retinoic acid-inducible gene-I (RIG)-like receptors that allow them to recognize several different types of Pathogen-associated molecular patterns (PAMP). In contrast to cDC1, these cells have the capacity to produce high amounts of IL-12, allowing them to activate Th1, Th17 and CD8⁺ T cells (64). Langerhans cells are skin-resident cells, specialized in production of pro-inflammatory cytokines while also presenting antigens to CD8⁺ T cells (64).

Several reports have shown that monocytes can migrate into tissues in response to local inflammation, rapidly acquiring key properties of DC such as expression of specific DC markers, present exogenous antigens to T cells and produce high amounts of inflammatory as well as anti-inflammatory cytokines (64). Monocytes differentiate into immature DC (iDC) specialized to engulf foreign antigens, resulting in antigens loaded into MHC molecules. Generation of mature DC (mDC) is achieved after activation of iDC with different antigens, mainly LPS, TNF- α or CD40L, leading to elevated expression of costimulatory molecules, migration to lymph nodes, antigen presentation to T cells and production of T cell-polarizing cytokines (68-70). Upon LPS recognition by TLR4, downstream adaptors are recruited into the intracellular domain of the receptor (Figure 1), such as the myeloid differentiation primary response gene 88 (MyD88), TIR domain-containing adaptor protein (TIRAP), TIR domain-containing adaptor inducing IFN- β (TRIF) and TRIF-related adaptor protein (TRAM). Different TLR use different combination of adaptor proteins to determine downstream signaling. TLR 1, 2, 5 and 6 mainly use the MyD88-dependent pathway, while TLR4 use both downstream signaling pathway. MyD88 activation leads to recruitment and activation of the IRAK and TRAF6 kinases, which activate I κ B kinase (IKK) and mitogen-activated protein kinase (MAPK) downstream signaling. IKK further phosphorylates I κ B which leads to its degradation and subsequent nuclear translocation of the NF- κ B transcription factor resulting in DC maturation by increasing the expression of costimulatory molecules and cytokines production (71-77). Activation of MAPK results in induction of the activator protein-1 (AP-1) transcription factor, which has also a role in the transcription of pro-inflammatory cytokines (71-73). Notably, MAPK regulates DC maturation through phosphorylation of NF- κ B. Accordingly, inhibition of p38 and JNK MAPK activation blocked the upregulation of the costimulatory molecules and the release of pro-inflammatory cytokines induced by LPS stimulation, resulting in a reduced capacity to prime T cells (74, 78, 79). In contrast, the

MyD88-independent pathway leads to activation of a different transcription factor interferon response factor (IRF), which is mainly involved in the type-1 interferon production (71-73).




Figure 1. TLR signaling pathway

Maturation of DC is typically achieved through stimulation with TLR agonists. Upon recognition of TLR ligands, signaling is driven in a MyD88-dependent or independent manner. MyD88 activation leads to recruitment of IRAK and TRAF6, followed by activation of MAPK and IKK. IKK phosphorylates IκB, leading to degradation of IκB and nuclear translocation of NF-κB. The p38, JNK and ERK MAPK activate AP-1 and NF-κB transcription factors. The MyD88-independent pathway is mediated through the TRIF/TRAM adaptor protein complex, which enhances IRF nuclear translocation. Finally, AP-1, NF-κB and IRF mediate the transcription of genes involved in costimulatory molecules expression and cytokine production. Figure modified after (71-73)

Due to the relevance of monocytes-derived DC *in vivo*, the development of protocols for *in vitro* generation of these cells has been used successfully to study DC biology. CD14⁺ monocytes can be isolated from donors and cultured in presence of GM-CSF and IL-4 for 6 days to obtain iDC. iDC generated by this method resemble the surface molecule expression and morphology observed *in vivo*. iDC are characterized by enhanced capacity to bind and

process antigens while expressing markers including group 1 CD1 (CD1a, CD1b and CD1c). Moreover, mDC retain expression of CD1 molecules and upregulate antigen-presenting and costimulatory molecules on their surface, including MHC class I and II, CD40, CD80, CD86 (64, 80-82).

3.3.2 DC antigen presentation and T cell polarization

DC have the capacity to efficiently present antigens to T cells. This process is in part dependent on the optimal expression of antigen-presenting molecules, including as MHC class I and II molecules. While MHC class I present peptide antigens to CD8⁺ cytotoxic T cells, MHC class II activates CD4⁺ T helper cells (67). MHC class II proteins include HLA-DR, HLA-DQ and HLA-DP, and are expressed on iDC and become highly up-regulated after iDC maturation. Efficient antigen presentation by DC correlates with the levels of these molecules on the surface (67, 83). Unlike MHC class I and II molecules which present peptides as antigens, members of the CD1 family present self- and foreign lipid antigens to T lymphocyte subsets. CD1 molecules are classified depended on their sequence homology into group 1 (CD1a-c), group 2 (CD1d) and group 3 (CD1e) (84, 85). While group 1 and 2 are expressed on the surface, group 3 is located intracellularly and might play a role in antigen-processing rather than presentation (84-86). Group 1 family members present lipid antigens to conventional T cells, while CD1d functions as antigen-presenting molecule to a specific T cell subset known as invariant Natural Killer T cells (iNKT) (87, 88).

In concert with signals triggered by peptide-MHC or lipid-CD1 complexes presentation, T cell activation requires a “second signal” triggered by costimulatory molecule interactions, mainly mediated by the expression of CD40, CD80 and CD86 on DC (89-91). Binding of CD80 and CD86 to their partner CD28 on T cells confers optimal production of IL-2, an important cytokine that promotes expansion and survival of T cells (90-92). However, the CD40/CD40L pathway also regulates T cell priming and differentiation. CD40 ligation on DC further increases the expression of costimulatory, adhesion and MHC molecules (89, 93). Consequently, DC produce an array of cytokines that direct CD4⁺T cell differentiation into one of the different subsets of effector T cells, such as T helper 1 (Th1), Th2, Th17 or regulatory T cells (Treg) (94). Th1, Th2 and Th17 subsets are defined based on the different cytokine combinations that they secrete. Th1 cells development is induced when IL-12 is

present, and play an important role in activating monocytes, macrophages and CD8⁺ T cells, via secretion of pro-inflammatory cytokines, including IFN- γ (95-97). In contrast, Th2 differentiation requires IL-4 and are involved in the recruitment of eosinophils, mast cells and basophils, while also promoting the activation of B cells (96). Th17 arise when the cytokine IL-6 and transforming growth factor (TGF- β) are present. Th17 cells secrete IL-17 and IL-22 and are known to be important in controlling fungal infections (98-100). Treg are crucial in the maintenance of peripheral tolerance, by the release of anti-inflammatory cytokines such as TGF- β and IL-10 (101). Treg cell population has been specially described through their expression of the transcription factor FOXP3, which is critical for the development and function of these cells (102, 103). DC promote tolerance by limiting effector T cells and promoting Treg differentiation in the periphery through various mechanisms including the production of immunosuppressive cytokines such as IL-10, IL-27 and TGF- β , and the expression of inhibitory molecules such as PD-L1, ILT2, ILT3 and ILT4 (104-107). In general terms, the interaction between PD-L1 and its receptor PD-1 inhibits immune cells activation via induction of apoptosis and T cell anergy (108). Moreover, engagement of the inhibitory receptors ILT2, ILT3 and ILT4 reduces the expression of costimulatory and antigen-presenting molecules, and inhibits the secretion of cytokines and chemokines in DC (109).

3.3.3 DC signaling and immunity against *Candida albicans*

DC play an important role in controlling mucocutaneous and systemic *C. albicans* infections (110-112). For *C. albicans*, dimorphism is a key trait involved in the interaction with DC. Ingestion of *C. albicans* yeast cells activates IL-12 production, promoting a priming of Th1 cells. In contrast, ingestion of hyphae leads to non-protective Th2 priming by suppressing IL-12 release and increasing IL-4 production (113). Moreover, immunization of mice with DC pulsed with *C. albicans* yeasts, but not hyphae, reduced the CFU observed in kidneys of *C. albicans* infected mice (113). While Th1 responses are known to be relevant during systemic infection, Th17 play an important role during mucocutaneous infection. Th17 cell activation by DC is induced by yeast cells in a process mainly dependent on the dectin-1/2-SYK-CARD9 signaling (99, 111, 114, 115).

DC effectively recognize *C. albicans* through the expression of a wide range of pattern recognition receptors (PRR), including the C-type lectin receptors (CLR) dectin-1, dectin-2 and cell-specific intracellular adhesion molecule-3-grabbing non integrin (DC-SIGN), as well as the mannose receptor (MR), Mincle, Galectin-3, TLR2, TLR4, TLR6, and complement receptor 3 (CR3). Altogether, these receptors efficiently recognize structural conserved moieties expressed on the cell wall of *C. albicans*, such as β -glucans, phospholipomannan, *N*-linked and *O*-linked mannans (116-122). However, the CLR dectin-1 and dectin-2 play an essential role in morphology-specific recognition of *C. albicans*. While dectin-1 recognizes β -glucan residues mainly expressed in yeast cells, dectin-2 recognizes mannose structures present mostly in hyphae forms (123). Although these receptors have certain differences in the downstream signaling compared to TLR, they also share common pathways including activation of NLRP inflammasomes, MAPK, NF- κ B and AP-1 transcription factors, resulting in enhanced the expression of costimulatory molecules and secretion of cytokines (Figure 2) (123-126).




Figure 2. Dectin-1 and dectin-2 signaling pathway

Recognition of *C. albicans* antigens by dectin-1 and dectin-2 leads to phosphorylation of the spleen tyrosine kinase (Syk). Syk phosphorylation leads to NF- κ B nuclear translocation by removing its inhibitor $\text{i}\kappa\text{B}$ through the IKK complex. Syk can also activate the PLC γ 2 signaling leading to PKC activation that controls ROS/NLRP3 inflammasome, AP-1 activation via MAPK phosphorylation and activation of CARD9/Bcl10/Malt1 complex. The Card9 complex regulates NF- κ B activity through the iKK complex. Resultant activation of NF- κ B and AP-1 leads to expression of cytokines and chemokines. Figure modified after (126).

3.3.4 Nuclear receptors and their role in DC maturation and immune response against fungal infections

3.3.4.1 Nuclear receptors

Nuclear receptors (NR) are transcription factors activated by lipid-soluble ligands, which regulate the expression of several genes involved in diverse processes, including differentiation, development, proliferation, metabolism and inflammation (127-130). NR transcriptional activity is achieved by a conserved structure consisting of regions with specific functions as depicted in figure 3. All NR contain an activation function region 1 (AF-1), a hypervariable N-terminal region which serves as a target for post-translational modifications, such as phosphorylation and SUMOylation, facilitating ligand-independent activation function. Furthermore, NR contain a highly conserved DNA-binding domain (DBD) which allows the binding to specific DNA sequences known as hormone response element. NR ligands bind to the C-terminal region containing the ligand-binding domain (LBD), which also mediates dimerization with another NR. Finally, the AF-2 conserved region located near the C-terminus regulates ligand-dependent activation of NR, by controlling the binding of coactivator and corepressor complexes (127, 128, 130, 131).



Figure 3. Nuclear receptor structure

The superfamily of NR is subdivided based on their ligands as “classical” endocrine receptors, orphan receptors and adopted orphan receptors (127, 128, 132). Classical receptors include the retinoic acid receptor (RAR), vitamin D receptor (VDR), thyroid hormone receptors (TR), which are activated by *all-trans* retinoic acid, 1 α 25-dihydroxyvitamin D, and thyroid hormones, respectively (127). In contrast, the ligands for orphan receptors are not known. However, several NR became “adopted” following the description of several ligands, including peroxisome proliferator activated receptor (PPAR), liver X receptor (LXR), farnesoid X receptor (FXR) and retinoid X receptor (RXR), activated mainly by metabolites such as fatty acids, oxysterols and bile acids (127, 128, 132).

One-third of NR form an obligatory heterodimeric partner with RXR and can be further classified in two groups depending on the ligand binding properties: permissive and non-permissive NR. While permissive NR, including PPAR and LXR can be activated either by their own ligands or RXR ligands, non-permissive NR, such as RAR, VDR and TR become only activated by their own agonists (127, 128). RXR heterodimers are located in the nucleus and bind to specific nucleotide sequences, known as hormone response elements, in the absence of ligands inhibiting transcriptional activation by the recruitment of corepressor molecules (Figure 4A) (127). However, the presence of ligands promotes dissociation of the corepressors and recruitment of coactivator complexes necessary for optimal transcriptional activity (Figure 4B). Alternatively, some ligands can trigger corepressor molecules recruitment, which blocks transcriptional properties of NR (Figure 4C). Finally, other NR suppress activation of different transcription factors in a process known as transrepression (Figure 4D). Particularly, SUMOylation of PPAR γ and LXR triggers their recruitment to promoters of inflammatory transcription factors, including NF- κ B and activator protein 1 (AP-1) and physically modulating their activation (127).




Figure 4. Nuclear receptor mechanism of action

Nuclear receptors and its partner, RXR, binds to specific DNA sequences and regulates the expression of several genes involved in multiple processes. Nuclear receptors activation is tightly regulated by the presence of its ligands. Liganded receptors can recruit co-repressors or co-activator complexes to enhance or inhibit gene expression. Alternatively, nuclear receptors might interfere with activation of other transcription factors in a process known as transrepression. Figure modified after (127).

3.3.4.1.1 Retinoid X receptor: RXR

Retinoic X receptor (RXR) is a unique member of NR because it forms heterodimers with several other NR, most of them requiring RXR as a mandatory partner for DNA binding and transcriptional activity. Thus, binding of ligands to permissive RXR heterodimers can potentially modify various pathways simultaneously, thereby modulating developmental, metabolic and immune processes (127, 133, 134). The most common ligand known is the vitamin A metabolite 9-cis retinoic acid (9-cis-RA) which is able to activate all three forms of RXR - RXR α , RXR β and RXR γ (135). Nevertheless, several other synthetic and natural agonists have been described (136).

The regulation of immune function by this receptor was shown in human monocytes-derived DC (137). Particularly, activation of RXR α regulates the differentiation process from monocytes to DC by altering the expression of costimulatory and antigen-presenting molecules, reducing IL-12 release, and increasing IL-10 production in response to LPS stimulation (137, 138). Furthermore, activation of RXR α *in vivo* reduced DC migration to draining lymph nodes due to inhibition of CCR7 expression (139).

3.3.4.1.2 Retinoic acid receptor: RAR

Retinoic acid receptor (RAR) mediates all its functions by recognition of the active metabolite of vitamin A, *all-trans* retinoic acid (ATRA), or synthetic ligands which can activate the three isoforms – RAR α , RAR β and RAR γ . RAR plays a crucial role in the regulation of immune function, highlighted by higher rates of infections observed in vitamin A deficient patients (127, 140-143). RAR forms a non-permissive heterodimer with RXR that binds to the retinoic acid response element, consisting of two direct repeats of PuGGTCA spaced by five bases (144). Similar to other NR, RAR can interact with several other pathways which highlight its relevance in the regulation of multiple processes (145-147).

Several *in vitro* and *in vivo* studies suggested an anti-inflammatory effect of RAR activation on innate immune cells (148-150). RAR activation by ATRA or synthetic agonists promotes the differentiation of M2 macrophages expressing anti-inflammatory cytokines, while suppressing TNF- α and nitric oxide production (148). Moreover, activation of RAR by ATRA leads to a repression in IL-12 release via inhibition of NF- κ B activation (149, 150).

RAR is also involved in modulating DC function. Particularly, DC derived from human cord blood and differentiated in presence of ATRA are characterized by a reduced IL-12 and increased IL-10 secretion (151). Furthermore, DC priming with ATRA enhances the release of TGF- β , which promotes expansion of Treg cells, highlighting the importance of ATRA as a metabolite with immune-suppressive potential (152, 153). Additionally, treatment of mouse splenic DC with ATRA promotes the expression of SOCS3, which attenuates the activation of p38 MAPK, and thereby, inhibits the release of IL-12, IL-6 and TNF- α (154). However, in presence of pro-inflammatory stimuli, ATRA treatment further increases the maturation of DC by activation of NF- κ B (155). Interestingly, activation of RAR showed to coordinate several genes during the transition from monocytes to DC. Particularly, group 1 of CD1 genes (CD1a, CD1b and CD1c) was downmodulated, while CD1d showed upregulation upon ATRA treatment. Moreover, upregulation of CD1d correlated with an elevated capacity to activate invariant Natural Killer T cells (iNKT) (156). Finally, ATRA could be used as an adjuvant due to the enhanced DC migration to draining lymph node in a murine model (157).

Activation of RAR α in immune cells modulates infection by pathogenic fungi. In particular, ATRA treatment diminished inflammation induced by monocytes infected with *C. albicans* by reducing the surface exposure of Dectin-1 and TLR2, together with a diminished

secretion of TNF- α , IL-6 and IL-12. The authors concluded that ATRA administration might be useful for prophylactic applications in septic patients infected with *C. albicans* (158).

3.3.4.1.3 Peroxisome proliferator activated receptors: PPAR

The family of peroxisome proliferator-activated receptors (PPAR) includes three isoforms: PPAR α , PPAR β/δ and PPAR γ (159). These isoforms differ among each other in their tissue expression. PPAR α is involved in fatty acid metabolism processes, including β -oxidation pathway, fatty acid catabolism, lipogenesis and ketone body synthesis (160). Thereby, PPAR α is expressed in metabolically active tissues, such as liver, heart, skeletal muscle, intestinal mucosa and brown adipose tissue (160, 161). Immune cells including monocytes, macrophages and lymphocytes also express this NR (162, 163). Although PPAR β/δ is ubiquitously expressed in all organs, it is highly present in the gastrointestinal tract (esophagus, liver, and intestines), kidneys and skeletal muscle (164). Similarly, PPAR γ is expressed in white and brown adipose tissue, the large intestine, spleen, and immune cells, most prominently in macrophages and DC (127, 160, 165).

PPAR have a large cavity binding site, which is approximately 3-4 larger than other NR (166). Thereby, PPAR are able to bind several natural and synthetic ligands. In particular, PPAR γ is activated by natural ligands such as unsaturated fatty acids, 15-hydroxy-eicosatetraenoic acid, 9- and 13-hydroxy-octadecadienoic acid, 15-deoxy Δ 12,14-prostaglandin J2 and prostaglandin J2 (160, 165). Among synthetic ligands are the insulin sensitizers Thiazolidinediones (such as englitazone, ciglitazone, pioglitazone, troglitazone and rosiglitazone), non-steroidal anti-inflammatory drugs (i.e. indomethacin, fenoprofen, flufenamic acid) and other PPAR γ modulators, so-called SPPARMs (Selective PPAR modulators) (165).

PPAR γ is known to influence the development and functionality of monocytes and macrophages (167-171). Activation of PPAR γ suppressed the production of pro-inflammatory cytokines, such as TNF, IL-6, IL-1 β , IL-12, iNOS and MMP-9 in these cells (167-170). Interestingly, PPAR γ also modulates macrophages migration in response to MCP-1, by suppressing CCR2 expression (171). PPAR γ inhibits inflammation by physically interacting with other transcription factors involved in pro-inflammatory pathways, including NF- κ B, AP-1, STAT-1 and NFAT, in a process known as transrepression. This process is

regulated by SUMOylation of PPAR γ , which allows PPAR γ to directly interact with the transcription factors, thereby preventing the binding to their response elements (172). Sequestration of coactivators, signaling through the IL-4 receptor, and altered activation of MAPK mediated by PPAR γ seems to play an important role in suppressing production of inflammatory cytokines (127, 173-175).

Furthermore, PPAR γ is well known to modulate DC functionality by altering the surface phenotype, cytokine secretion and T cell activation capacity. Particularly, treatment of DC with PPAR γ agonists dampens the release of pro-inflammatory cytokines and chemokines, including TNF- α , IL-12, IL-5, IL-6, IP-10 (CXCL10) and RANTES (CCL5) (176, 177). Moreover, PPAR γ activation enhances CD86 while downmodulates CD80 expression (87). However, the most important effect was observed in CD1 molecule expression pattern. While CD1a is downregulated, CD1d expression increases in PPAR γ -instructed DC (87). Mechanistically, PPAR γ modulates CD1d expression indirectly by enhancing the activity of aldehyde dehydrogenase (ALDH1A, RALDH) which is involved in ATRA production. ATRA binds to RAR and modulates the expression of CD1d (156). Furthermore, PPAR γ activation increases the expression of cathepsin D, a lysosomal aspartyl protease, which improves lipid antigen processing and loading to the CD1d complex (178).

PPAR γ may play either a beneficial or detrimental role during infection, mainly dependent on the pathogen involved (179-181). Majer *et al.* showed that administration of the synthetic PPAR γ agonist pioglitazone improves viability of mice infected by *C. albicans* by suppressing neutrophils infiltration into the kidneys of infected mice (182). Furthermore, PPAR γ activation increases the expression of dectin-1 and MR in macrophages. Accordingly, macrophages produce more ROS and have a higher capacity to uptake and kill *C. albicans* (183-185).

3.3.4.1.4 Liver X receptor: LXR

Liver X receptors (LXR) – namely LXR α and LXR β - have a large hydrophobic cavity that enables the binding of cholesterol and several oxysterols. Thus, this NR play a key role in modulating metabolism and cholesterol homeostasis (186). LXR is highly expressed in the liver, intestine, adipose tissue, lung, the adrenal glands, kidney, lymphocytes, monocytes, macrophages and DC (127, 186). LXR forms a permissive heterodimer with RXR, and binds

to specific DNA sequences known as LXR-responsive elements (LXRE), consisting in direct repeats of AGTCA separated by 4 nucleotides (187).

Most of the studies focused on the role of LXR in modulating lipid metabolism. However, LXR activity also regulates immune responses, especially in macrophages and DC (188-192). LXR exerts anti-inflammatory effects in macrophages by inhibiting the expression of several pro-inflammatory molecules, such as IL-6, IL-1 β , MCP-1, MCP-3, iNOS, and cyclooxygenase 2 in response to bacterial or LPS stimulation (193, 194). Similarly as PPAR γ , LXR α can suppress inflammatory signaling by transrepression which requires SUMOylation of the receptor (195). Interestingly, these effects were only observed in short-term LXR stimulation. Long-term activation of LXR in human macrophages potentiates LPS responses, by increasing TLR4 expression, activation of MAPK and ROS production (196).

The role of LXR in DC biology is conflicting. Geyeregger *et al.* showed that treatment of monocytes with a synthetic LXR ligand impairs DC differentiation and maturation by reducing IL-12 and increasing IL-10 secretion (190). Moreover, these cells have a lower capacity to activate T-cell, in a process dependent on the reduction of the actin-bundling protein fascin (190). However, comprehensive transcriptional and functional analysis showed that differentiation of human DC in presence of LXR ligands promotes their capacity to activate CD4⁺ T cells by increasing the expression of costimulatory molecules and pro-inflammatory cytokines production (IL-12, TNF- α , IL-6 and IL-8), in a process dependent on NF- κ B activation (191). Interestingly, LXR has a crucial role in DC migration during inflammatory conditions due to the regulation of the ectoenzyme CD38 (192, 197).

Several reports showed that LXR play an important role in modulating the immune response against pathogenic bacteria, viruses and parasites (188, 198-202). However, only one report has addressed the role of LXR during infection by pathogenic fungi. Bobryshev *et al.* demonstrated that macrophages infected with *Chlamydia pneumoniae* showed reduction in LXR α expression, which correlated with higher formation of foam cells in atherosclerotic lesions (203).

In summary, NR modulate differentiation and maturation of DC. Furthermore, these receptors are involved in the interaction with pathogenic microorganisms. Nevertheless, only few reports have addressed the role of these transcription factors during fungal infections. Moreover, no study so far has evaluated the potential of microbial products produced by pathogenic fungi in modulating immune responses through NR activation.

4 Aims of the work

Farnesol, the first quorum sensing system described in eukaryotes, modulates the hyphae to yeast transition in the filamentous fungus *C. albicans*. However, farnesol also acts as a virulence factor during *C. albicans* infection, by altering immune cells function. Particularly, farnesol impairs dendritic cell differentiation and maturation process from monocytes. Nevertheless, the molecular mechanisms by which farnesol alters dendritic cell functionality remain elusive. In this work, the elucidation of signaling pathways activated by farnesol in dendritic cells was investigated.

The specific aims of this work were:

- Define molecular pathways modulated by farnesol through the available microarray data of dendritic cells differentiated in presence of farnesol
- Characterize the contribution of nuclear receptors to modulation of dendritic cell maturation by farnesol
- Analyze downstream inflammatory signaling pathways activated by farnesol during the maturation process of monocytes to dendritic cells
- Evaluate the impact of farnesol treatment on the ability of dendritic cells to activate different T cells subsets

5 Experimental procedures

5.1 Reagent preparation

Trans-trans, FOH was obtained as a 4M stock solution (Sigma-Aldrich) and then diluted in 100% methanol, followed by serial dilutions in RPMI + 5% FBS to the working concentrations of 50 μ M and 100 μ M. Concentrations of FOH were selected based on the observation that *C. albicans* can release up to 55 μ M FOH in *in vitro* culture, and concentrations of 10-250 μ M have been shown to modulate morphology transition in *C. albicans* (204, 205). The following antagonists for nuclear receptors were obtained from Tocris Bioscience: GW9662 (PPAR γ antagonist, 10 μ M), AGN193109 (RAR α antagonist, 10 μ M), GSK2033 (LXR α antagonist, 1 μ M). Nuclear receptor agonists were: Rosiglitazone (PPAR γ agonist, 5 μ M, Enzo Life Sciences), AM580 (RAR α agonist, 100 nM, Sigma-Aldrich) and GW3965 (LXR α agonist, 1 μ M, Tocris Bioscience). Methanol was used as a solvent control indicated as mock-treated DC. Lipopolysaccharide (LPS, Sigma-Aldrich), peptidoglycan (Sigma-Aldrich), flagellin (Sigma-Aldrich), zymosan (InvivoGen) and Pam3CSK4 (R&D Systems) were diluted in water before application. α -galactosylceramide (α -GalCer) was obtained from AbCam and diluted in DMSO.

5.2 Dendritic cell generation and maturation

Human monocytes were isolated from buffy coats of healthy volunteers (provided by the Transfusion Medicine Department of the University Hospital Jena). First, Peripheral Blood Mononuclear Cells (PBMC) were obtained by density gradient centrifugation using BIOCOLL (Biochrom AG), and monocytes were positively selected by using anti-CD14-conjugated beads via a magnetic cell sorting system (MACS) (Miltenyi Biotec). Freshly isolated monocytes were resuspended into 6-well plates at a density of 2×10^6 cells per well and cultured in RPMI-1640 (Biochrom AG) with 10mM L-glutamine supplemented with 10% heat inactivated FBS (Biochrom AG), 800 U/ml GM-CSF (Leukine® Sargamostim) and 1000 U/ml IL-4 (Miltenyi Biotec). Cells were cultured for 6 days at 37°C and 5% CO₂ and half of media was changed at day 3. Ligands or solvent control were added to the cell culture starting from the first day. NR antagonists were added 1 hour prior FOH or solvent stimulation. iDC differentiation was confirmed by CD1a positive and CD14 negative staining.

To obtain matured dendritic cells (mDC), iDC were treated with 10 ng/ml LPS (Sigma-Aldrich) for 24 hours. To block NF- κ B and MAPK activation, iDC were pre-incubated for 30 min prior LPS stimulation with the following inhibitors: SC75741 (NF- κ B inhibitor, 5 μ M, Cayman), SP600125 (JNK inhibitor, 10 μ M, InvivoGen), SB203580 (p38 inhibitor, 20 μ M, InvivoGen), and FR180204 (ERK1/2 inhibitor, 10 μ M, Tocris). To evaluate Toll-like receptor and dectin-1 activation, iDC were stimulated for 24 hours with the following ligands: Pam3CSK4 (TLR 1/2 ligand, 10 μ g/ml, R&D systems), peptidoglycan (TLR 2/6 ligand, 10 μ g/ml, Sigma), flagellin (TLR 5 ligand, 100 ng/ml, Sigma) and zymosan (dectin-1 ligand, 100 μ g/ml, InvivoGen) (Figure 5).




Figure 5. Generation of monocytes-derived DC

5.3 *Candida albicans* cultivation

Wild-type *C. albicans* strain SC5314 was maintained on YPD agar. For experiments, cells were transferred to M199 medium, pH 4 (9.8 g/l M199, 35.7 g/l HEPES, 2.2 g/l NaHCO₃), and cultured at 30°C overnight to stationary phase. Germ tubes were induced by culturing in M199 medium, pH 8, for one hour at 37°C. Germ tubes were washed with HBSS and killed by adding 0.1 % Thimerosal (Sigma-Aldrich) for one hour at 37°C. MOI of 1 was used for confrontation assays.

5.4 Flow cytometry

All flow cytometric analyses were performed using the FACSCanto II (BD Bioscience) and the data was analyzed with FlowJo 7.6.4 software.

5.4.1 Viability determination

For determination of viability, cells were harvested and stained with Fixable Viability Stain V450 (BD Bioscience) for 15 min at room temperature prior cell surface staining. The percentage of viable cells (negative staining) was measured.

5.4.2 Characterization of cell surface markers

Phenotypic characterization of DC, iNKT, Th1 and Treg cells was performed by flow cytometry using fluorochrome-conjugated antibodies. 100 µl of cell suspension was stained with the following antibodies (See table 1). Isotype controls and fluorescence minus one (FMO) control were used for proper gating. After 20 min of staining, cells were resuspended in CellWash (BD Bioscience) and analyzed immediately on the flow cytometer.

Antibody/origin	Isotype/origin	Concentration
FITC anti-human CD14	Mouse IgG2a/Biolegend	100 µg/ml
PE anti-human CD1a	Mouse IgG1/Biolegend	50 µg/ml
PE anti-human CD80	Mouse IgG1/Biolegend	100 µg/ml
V450 anti-human CD86	Mouse IgG1/BD Bioscience	100 µg/ml
FITC anti-human CD40	Mouse IgG1/Biolegend	50 µg/ml
APC anti-human CD1d	Mouse IgG/Biolegend	100 µg/ml
PerCP anti-human HLA-DR	Mouse IgG2a/Biolegend	100 µg/ml
PerCp anti-human CD3	Mouse IgG1/Biolegend	50 µg/ml
PacificBlue anti-human CD4	Mouse IgG1/Biolegend	100 µg/ml
PE anti-human CD25	Mouse IgG1/Biolegend	25 µg/ml
APC anti-human CD85k (ILT3)	Mouse IgG1/Biolegend	100 µg/ml
FITC anti-human CD85j (ILT2)	Mouse IgG2b/Biolegend	100 µg/ml
PE anti-human CD85d (ILT4)	Mouse IgG2a/Biolegend	200 µg/ml
Brilliant Violet 421 anti-human CD274 (PD-L1)	Mouse IgG2b/Biolegend	50 µg/ml

PE anti-human TLR4	Mouse IgG2a/Biolegend	100 µg/ml
FITC anti-human 6B11 (anti-Vα24-Jα18)	Mouse IgG1/Biolegend	200 µg/ml

Table 1. Antibodies used for cell surface molecules and intracellular staining analysis

5.5 Invariant Natural Killer T cells expansion

Autologous T cells were collected by positive selection of CD3⁺ cells with microbeads (Miltenyi Biotec) and stored in DMSO + 90%FBS at -80°C until used in co-culture experiments. Mature dendritic cells were generated from primary monocytes as described above (Section 5.2). At day 7, mDC were treated with 100 ng/ml of α-GalCer for 24 hours to obtain α-GalCer-loaded mDC and were co-cultured with autologous T cells in a 1:10 ratio (1x10⁵ mDC : 1x10⁶ T cells) in 24-well plates. 24 hours later, 100 U/ml of recombinant human IL-2 (ImmunoTools) was added to induce T cell expansion. After 7 days, iNKT expansion was addressed by quantifying CD3⁺Vα24Jα18⁺ cells by flow cytometry. For cytokine reconstitution assays, recombinant human IL-12 (10 ng/mL) (ImmunoTools) and a mouse anti-human IL-10 blocking antibody (10 µg/mL) or a mouse IgG_{2B} isotype control (both obtained from R&D systems) were added from the first day of co-culture. At least 5x10⁵ events were acquired in the flow cytometer.

5.6 Intracellular staining

After 7 days of mDC – T cell co-culture, cells were collected and re-stimulated for 6 hours with PMA (50 ng/mL) and ionomycin (1 µg/mL), in presence of brefeldin A (5 µg/mL) and monensin (5 µg/mL) (Sigma-Aldrich). Intracellular staining of IFN-γ with FITC anti-human IFN-γ antibody (Biolegend, concentration: 25 µg/ml) and intracellular fixation and permeabilization buffer set according to the manufacturer's protocol (eBioscience). To analyze Treg expansion, cells were fixed and permeabilized with the eBioscience FOXP3/Transcription factor staining set, followed by staining with FITC anti-human FOXP3 (Biolegend, concentration: 80 µg/ml). CD3⁺IFN-γ⁺ and CD3⁺CD25⁺FOXP3⁺ cells were quantified by flow cytometry.

5.7 Phosphoflow

For experiments analyzing NF- κ B and p38 MAPK phosphorylation, 5×10^5 DC/well were seeded into 24-well plates and stimulated with 10 ng/ml LPS for 1 hour, 37°C and 5% CO₂. To stop the reaction, cells were fixed with intracellular fixation buffer (eBioscience) for 30 min at RT and permeabilized with 90% freezing methanol overnight. Cells were stained by using primary antibodies to phospho-NF- κ B p65 (Ser536; clone 93H1) or phospho-p38 MAPK (Thr180/Tyr182; clone 12F8) (Cell signaling) for 1 hour in the dark at RT followed by staining with Dylight649-conjugated donkey anti-rabbit or Dylight488-conjugated donkey anti-mouse (Jackson ImmunoResearch) antibodies for 15 min at RT. PBS containing 0,5% BSA was used for washing and intracellular staining. Cells were analyzed in the flow cytometer.

5.8 Cytokine measurement

iDC were incubated in presence of either TLR and dectin-1 ligands for 24 hours, and then supernatants were collected and stored at -80 °C. Cytokine concentrations were analyzed with the ProcartaPlex immunoassay from ThermoFischer Scientific according to the manufacturer's instructions.

5.9 RNA isolation and quantitative real-time PCR

Preparation of whole cell RNA was carried out with the RNeasy Plus Mini Kit (Qiagen) according to the manufacturer's instructions. NanoDrop (ThermoFisher) was used for the quantification of the RNA preparations. cDNA was generated from 25 ng of total RNA with the Precision nanoScript 2 Reverse transcription kit (Primer Design) and qRT-PCR was performed using the PrecisionPLUS MasterMix premixed with SYBRgreen kit (Primer Design) and Quantitect Primer Assay oligonucleotides (Qiagen) on a Stratagene Mx3500P cyclor. Gene expression was normalized to the TBP reference gene and relative expression was calculated using the $2^{-\Delta\Delta CT}$ method.

5.10 Statistical analysis

Data were analyzed with GraphPrism software version 6. Shapiro-Wilk normality test was used to assess Gaussian distribution. Comparison among groups was performed using a one-way ANOVA with Turkey's posttest for multiple comparisons. Otherwise, Kruskal-Wallis was used as a non-parametric test. Two-tailed unpaired Student's t test was applied for comparison between two groups. $P < 0.05$ was considered as significant and marked in graphs using the following designations: * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$.

6 Results

6.1 Viability of monocyte-differentiated DC in presence of farnesol

Farnesol (FOH) treatment inhibits proliferation while inducing apoptosis in several type of cells (51-57). Thus, Leonhardt *et al.* performed a viability analysis of iDC with Annexin-V and PI staining and showed that FOH treatment during the differentiation process did not influence the viability of iDC when used in concentrations up to 50 μ M, while retaining its effects on DC differentiation (56). To confirm this observation, viability of iDC generated in presence of FOH was investigated with the use of the fixable viability stain from BD. This dye binds to amines which are accessible in permeable cell membranes present in late-apoptotic/necrotic cells. As shown in figure 6, no differences could be observed in cell viability in iDC differentiated in presence of the solvent control or FOH (viable cells: mock-treated, 94 ± 5 %; with 100 μ M FOH, 89 ± 7 %) when compared to untreated cells (94 ± 4 %). Since we observed no differences in viability for FOH concentrations 50 μ M and 100 μ M, subsequent experiments were performed using both FOH concentrations.




Figure 6. Viability of iDC differentiated in presence of FOH

Monocytes were differentiated into iDC in absence or presence of FOH (50 μ M and 100 μ M) and 1% methanol (mock-treated). Quantitative data of viable cells are shown as means \pm SD from 4 independent experiments using cells from different donors.

6.2 Farnesol-differentiated DC have altered activation in response to distinct stimulation

Previous experiments performed by Leonhardt *et al.* showed that FOH impairs DC differentiation from monocytes by altering the surface molecule expression and the secretion of Th1-inducing cytokines (56). However, all the demonstrated effects of FOH on mDC so far were determined after stimulation of Toll-like receptor 4 (TLR4) signaling by LPS. Thus, FOH could be influencing DC maturation by changing the expression pattern of this PRR. Nevertheless, the presence of FOH throughout the differentiation and maturation process did not modify TLR4 surface expression (Figure 7). To get a broader view regarding the impact of FOH, DC were differentiated and stimulated with inactivated *C. albicans* germ-tubes and several agonists for Toll-like receptors and dectin-1. Similar to LPS stimulation, the expression of costimulatory CD40, CD80, CD86, and the MHC class II molecule HLA-DR were modulated when cells were exposed to FOH. Differentiation in presence of FOH impairs the expression of CD40 and CD80 after every stimulation tested. Moreover, CD86 surface exposure after LPS stimulation was reduced in FOH-instructed DC compared to the mock-treated cells (DC with 100 μ M FOH, median of 2865 ± 515 ; mock-treated DC, median of 4589 ± 867 , $P < 0.01$). Interestingly, CD86 surface expression was enhanced in FOH-differentiated DC when stimulated with Pam3CSK4 (TLR 1/2 agonist) and flagellin (TLR 5 agonists) (DC with 100 μ M FOH: Pam3CSK4, median of 2286 ± 292 , $P < 0.01$; flagellin, median of 2135 ± 265 , $P < 0.001$) compared to mock-treated cells (mock-treated DC: Pam3CSK4, median of 1245 ± 547 ; flagellin, median of 819 ± 605). Furthermore, the MHC class II HLA-DR molecule was reduced in FOH-differentiated DC after stimulation with *C. albicans* (DC with 100 μ M FOH: median of 1186 ± 585 , mock-treated DC: median of 2410 ± 761 , $P < 0.05$) (Figure 8).

In addition, the release of cytokines and chemokines was altered in FOH-differentiated DC in response to Toll-like receptor and dectin-1 ligands (Figure 9). Production of pro-inflammatory IL-8 and TNF- α was enhanced in FOH-differentiated DC when treated with Pam3CSK4 and flagellin compared to mock-treated cells. Furthermore, the release of the chemokine MCP-1 (CCL2) and the anti-inflammatory IL-1 receptor antagonist (IL-1RA) was increased in FOH-differentiated DC in response to every tested stimulus. Similarly, the secretion of anti-inflammatory IL-10 was enhanced in FOH-differentiated DC when treated

with Pam3CSK4 and flagellin (DC with 100 μ M FOH: Pam3CSK4, 81 ± 72 pg/ml, $P < 0.001$; flagellin, 128 ± 124 pg/ml, $P < 0.05$) compared to mock-treated cells (mock-treated DC: Pam3CSK4, 13 ± 12 pg/ml; flagellin, 10 ± 6 pg/ml). Interestingly, the Th1-inducing cytokine IL-12 was reduced in FOH-differentiated DC stimulated with Toll-like receptors and dectin-1 ligands (DC with 100 μ M FOH: LPS, 120 ± 174 pg/ml; $P < 0.001$; peptidoglycan, 11 ± 16 pg/ml; $P < 0.001$) compared to mock-differentiated DC (mock-treated DC: LPS, 3840 ± 1924 pg/ml; peptidoglycan, 2958 ± 1698 pg/ml). Altogether, these results show that FOH alters the differentiation and maturation of DC by modifying the surface expression of key molecules and the release of cytokines and chemokines in response to several stimuli, indicating that FOH-induced changes are independent of any specific PRR and FOH affects downstream signaling cascades.




Figure 7. TLR4 expression in FOH-differentiated iDC and mDC

Monocytes were differentiated into iDC in the presence or absence of FOH and further stimulated with LPS for 24 hours. TLR4 median surface expression was quantified by flow cytometry. Quantitative data shown represent means \pm SD from 4 independent experiments with cells isolated from different donors.




Figure 8. CD40, CD80, CD86 and HLA-DR expression in FOH-differentiated DC activated with distinct stimuli

Monocytes were differentiated into iDC in the presence or absence of FOH and further stimulated with either LPS (TLR4 agonist), Pam3CSK4 (TLR1/2 agonist), peptidoglycan (TLR2/6 agonist), flagellin (TLR5 agonist), Zymosan (Dectin-1 agonist) and *C. albicans* germ-tubes for 24 hours. For each surface marker, median surface expression was quantified by flow cytometry. Quantitative data shown represent means \pm SD from 4 independent experiments with cells isolated from different donors (* $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$).




Figure 9. Cytokines and chemokines secretion of FOH-differentiated DC in response to distinct stimuli

Monocytes were differentiated into iDC in the presence or absence of FOH and further stimulated with either LPS (TLR4 agonist), Pam3CSK4 (TLR1/2 agonist), peptidoglycan (TLR2/6 agonist), flagellin (TLR5 agonist), Zymosan (Dectin-1 agonist), inactivated *C. albicans* for 24 hours. Cytokine concentrations were determined by multiplex assay. Data are means \pm SD of 4 independent experiments using cells isolated from different donors (* P < 0.05; ** P < 0.01; *** P < 0.001).

6.3 Molecular mechanisms modulated by farnesol on DC differentiation and maturation

6.3.1 Differentiation of DC in presence of farnesol increased the expression of genes involved in nuclear receptor activity

One of the most prominent effects in DC differentiated in presence of FOH was observed for the expression of CD1 transmembrane glycoproteins (56). FOH treatment shifts the expression pattern of CD1 molecules. While CD1a was not up-regulated in iDC and mDC, CD1d surface exposure was increased (Figure 10A). Consequently, the CD1a/CD1d ratio was reduced in FOH-differentiated DC (iDC with 100 μ M FOH, ratio of 0.3 ± 0.1) compared to the mock-treated cells (mock-treated iDC: ratio of 106 ± 51 , $P < 0.001$) (Figure 10B). Since expression of CD1 molecules in DC is known to be tightly regulated by the nuclear receptors PPAR γ and RAR α (87, 156), screening of NR and their target genes was performed from the transcriptional data available from Leonhardt *et al.* (56). Several genes linked to NR activation, including target genes for PPAR γ , RAR α , and LXR α were upregulated during DC differentiation in presence of FOH (Figure 11A). Quantitative real time PCR was used to validate the expression of NR and selected target genes. In accordance with the microarray data, FOH induced the expression of PPAR γ and its *bona fide* target gene *FABP4* as well as LXR α and its target gene *APOC1*. In addition, the RAR α target gene *TGM2* was upregulated in FOH-differentiated DC. Nevertheless, the expression of the receptor did not change during differentiation with FOH. The effects of FOH was concentration-dependent and a higher expression of these genes was observed in cells differentiated with 100 μ M FOH (Figure 11B).

These results confirmed FOH effects on CD1 molecules - while CD1a showed reduced expression, CD1d surface exposure was enhanced. Furthermore, FOH-differentiated DC have a higher expression of NR and their target genes.




Figure 10. CD1 expression of DC differentiated in presence of FOH

Surface CD1 molecules expression of iDC and mDC was analyzed by flow cytometry. (A) Representative histogram after 6 days of differentiation (iDC) and following LPS stimulation for 24 hours (mDC). (B) Data shows the ratio between measured expression of CD1a and CD1d on DC differentiated with FOH (50 µM and 100 µM), compared to mock-treated DC. The bars show means \pm SD from at least 4 independent experiments. (***) $P < 0,001$, compared to mock-treated DC)




Figure 11. Expression of nuclear receptors and their target genes in DC-differentiated in presence of FOH.

(A) Microarray data of immature and mature DC differentiated in absence or presence of FOH (50 μ M) was obtained from Leonhardt *et al.* and heat map of representative genes was generated. All displayed genes have a P value < 0.05 (56). (B) Confirmation of microarray results was performed by qRT-PCR. RNA was collected from iDC differentiated in presence or absence of FOH as described in experimental procedures. Gene expression was normalized to the TBP reference gene and relative expression was calculated using the $2^{-\Delta\Delta CT}$ method. Data are means \pm SD from 3 independent experiments using cells from different donors ($*P < 0.05$; $**P < 0.01$; $***P < 0.001$, compared to mock-treated DC).

6.3.1.1 Activation of PPAR γ and RAR α leads to a DC phenotype comparable to farnesol-differentiated DC

To evaluate the role of NR activation in the immunophenotypic changes induced by FOH, cells were exposed to specific NR agonists and antagonists throughout the process of DC differentiation and analyzed for the surface expression of costimulatory and antigen-presenting molecules.

Previous reports have shown that activation of NR induce a similar immunophenotype to FOH-differentiated DC (87, 156). In agreement with this, the presence of FOH, RSG (PPAR γ agonist) and AM (RAR α agonist) impaired the expression of CD1a after the differentiation of monocytes into immature DC (iDC) and after 24 hours of LPS stimulation (mature DC, mDC) compared to mock-treated DC (mock-treated iDC, median of 20737 ± 11256 , iDC with $100 \mu\text{M}$ FOH, median of 144 ± 62 , $P < 0.001$; iDC with RSG, median of 157 ± 38 , $P < 0.05$; iDC with AM, median of 274 ± 45 , $P < 0.001$) (Figure 12). Interestingly, expression of CD1d was elevated in iDC and mDC in presence of FOH, RSG and AM compared to mock-treated cells (mock-treated iDC, median of 255 ± 51 , iDC with $100 \mu\text{M}$ FOH, median of 638 ± 254 , $P < 0.01$; iDC with RSG, median of 496 ± 85 , $P < 0.05$; iDC with AM, median of 961 ± 393 , $P < 0.001$). No effect was observed when cells were treated with LXR α agonist GW3965. Moreover, the expression of the costimulatory molecule CD80 was diminished in iDC and mDC treated with FOH, PPAR γ and RAR α agonists compared to mock-treated cells (mock-treated mDC, median of 1565 ± 390 , mDC with FOH $100\mu\text{M}$, median of 725 ± 212 , $P < 0.001$; mDC with RSG, median of 951 ± 370 , $P < 0.05$; mDC with AM, median of 819 ± 123 , $P < 0.01$). In contrast, the expression of the costimulatory molecule CD86 was increased in iDC differentiated with FOH and agonists for PPAR γ and RAR α compared to mock-treated cells (mock-treated iDC, median of 139 ± 30 , iDC with FOH $100\mu\text{M}$, median of 732 ± 284 , $P < 0.05$; iDC with RSG, median of 2334 ± 1265 , $P < 0.001$; iDC with AM, median of 1181 ± 352 , $P < 0.01$). No differences in CD86 expression were observed after LPS stimulation. These results showed that differentiation of DC in the presence of PPAR γ and RAR α agonists lead to a similar immunophenotype of DC compared to FOH-differentiated cells.




Figure 12. Surface molecules expression on DC differentiated in presence of farnesol and nuclear receptors agonists

Monocytes were differentiated into DC in the presence or absence of 50 μ M or 100 μ M FOH and nuclear receptors agonists. For each surface marker, median surface expression after 6 days of differentiation (iDC) and 24 hours stimulation with LPS (mDC) was quantified by flow cytometry. Quantitative data shown represent means \pm SD from at least 3 independent experiments with cells isolated from different donors (* P < 0.05; ** P < 0.01; *** P < 0.001, compared to mock-treated DC). RSG: PPAR γ agonist Rosiglitazone; AM: RAR α agonist AM580; GW: LXR α agonist GW3965.

6.3.1.2 Nuclear receptor inhibitors have no significant effect on dendritic cell viability

DC were differentiated from monocytes in presence of previously published concentration of NR antagonists and viability was determined by flow cytometry. As shown in the figure 13, the presence or absence of FOH and NR antagonists during the differentiation process did not alter DC viability. Thus, subsequent experiments were performed by using these concentrations of FOH and NR antagonists.




Figure 13. Viability of DC differentiated with nuclear receptor antagonists

Discrimination of viable from non-viable cells was evaluated by flow cytometry. Dead cells were identified by positive staining with the Fixable Viability Stain V450. Data are means \pm SD from 3 independent experiments using cells isolated from different donors. PPAR γ antagonist: GW9662, RAR α antagonist: AGN193109, LXR α antagonist: GSK2033, RXR antagonist: HX531.

6.3.1.3 Farnesol modulates DC phenotype partially through activation of the PPAR γ and RAR α signaling pathways

DC generated in the presence of FOH were characterized by markedly lower surface expression of CD1a, but an increased expression of CD1d (Figures 10 and 12). To evaluate the role of nuclear receptor activation in the immunophenotypic changes induced by FOH, cells were exposed to various nuclear receptors antagonists during DC differentiation and analyzed for their surface expression of costimulatory and antigen-presenting molecules. The combination of FOH and NR antagonists were not enough to recover CD1a expression (Figure 14). In contrast, blocking the activity of PPAR γ with the PPAR γ antagonist GW9662 in iDC differentiated with 50 μ M and 100 μ M FOH returned CD1d expression to basal levels (mock-treated iDC, median of 266 ± 70 ; iDC with 50 μ M FOH, median of 481 ± 145 ; iDC with 50 μ M FOH and PPAR γ antagonist, median of 265 ± 53). A similar return to basal CD1d expression levels DC was observed in the presence of RAR α antagonist AGN193109 (mock-treated iDC, median of 289 ± 71 ; iDC with 50 μ M FOH, median of 474 ± 149 ; iDC with 50 μ M FOH and RAR α antagonist, median of 243 ± 64). These effects were also maintained during the maturation process from iDC to mDC, suggesting that the increased CD1d surface exposure induced by FOH is regulated by a common pathway that involves both receptors. The use of LXR α antagonist affected neither CD1a nor CD1d surface expression during FOH treatment. However, blocking LXR α receptor activity downregulated CD1a expression (mock-treated iDC, median of 12731 ± 6598 ; with LXR α antagonist, median of 2845 ± 2469) and promoted the surface expression of CD1d (mock-treated iDC, median of 259 ± 85 ; with LXR α antagonist, median of 581 ± 203) in mock-treated control cells comparable to the effects induced by FOH (iDC with 50 μ M FOH: CD1a, median of 255 ± 98 ; CD1d, median of 525 ± 227). To confirm the role of NR in the regulation of CD1 molecules by FOH, cells were differentiated in presence of an antagonist for RXR α , which serve as a partner for these receptors and are known to be essential for their functionality (127). Downmodulation of CD1a expression by FOH was not restored in presence of RXR α antagonist (Figure 15). Nevertheless, CD1d upregulation in FOH-instructed DC was blocked when the antagonist was present after the differentiation process (iDC with 100 μ M FOH, median of 400 ± 81 ; with 100 μ M FOH and RXR α antagonist, median of 241 ± 30 , $P < 0.05$), reaching levels observed in mock-treated cells (mock-treated iDC, median of 202 ± 53).




Figure 14. CD1 molecule expression induced by FOH in presence of nuclear receptor antagonists

Monocytes were differentiated into DC in the presence of either 50 μ M or 100 μ M FOH or mock treatment and the nuclear receptors antagonists indicated. For both surface markers, median surface expression after 6 days of differentiation (iDC) and 24 hours stimulation with LPS (mDC) was quantified by flow cytometry. Quantitative data shown represent means \pm SD from at least 5 independent experiments with cells isolated from different donors (* P < 0.05; ** P < 0.01; *** P < 0.001). PPAR γ antagonist: GW9662, RAR α antagonist: AGN193109, LXR α antagonist: GSK2033, RXR antagonist: HX531.




Figure 15. CD1 molecule expression induced by FOH in presence of RXRα antagonist

Monocytes were differentiated into DC in the presence of either 50 μM or 100 μM FOH or mock treatment and the RXR antagonist. For both surface markers, median surface expression after 6 days of differentiation (iDC) and 24 hours stimulation with LPS (mDC) was quantified by flow cytometry. Quantitative data shown represent means \pm SD from 4 independent experiments with cells isolated from different donors (* P < 0.05; ** P < 0.01; *** P < 0.001). RXRα antagonist: HX531.

To fully characterize the role of NR on the impact of FOH over DC differentiation and maturation, the expression of costimulatory and antigen-presenting molecules, which play a key role in DC functionality, was analyzed by flow cytometry. CD80 and CD40 expression was reduced in FOH-differentiated iDC and mDC (Figure 16). Nevertheless, the presence of PPAR γ antagonist partially blocked CD80 downmodulation by lower FOH concentration in iDC (iDC with 50 μ M FOH, median of 276 ± 65 ; iDC with 50 μ M FOH and PPAR γ antagonist, median of 454 ± 120 , $P < 0.05$) and reached levels similar to mock-treated DC (median of 385 ± 140). In contrast, CD86 was upregulated in cells differentiated with FOH and the presence of PPAR γ antagonist blocked this effect (mock-treated iDC, median of 202 ± 79 , iDC with 50 μ M FOH, median of 505 ± 207 , iDC with 50 μ M FOH and RAR α antagonist, median of 242 ± 80 , $P < 0.05$). Similarly, the presence of RAR α antagonist blocked the upregulation of CD86 induced by FOH treatment (Figure 17). Interestingly, CD86 expression was partially down-modulated in presence of FOH after LPS stimulation and this effect was blocked during treatment with PPAR γ antagonist. Finally, no significant differences were observed in HLA-DR expression after differentiation in presence of FOH. Nevertheless, HLA-DR expression was diminished in FOH-differentiated DC after LPS stimulation with no significant effects observed in presence of NR antagonists (mock-treated mDC: median of 3669 ± 213 , mDC with 100 μ M FOH, median of 2360 ± 994 , $P < 0.05$).

Altogether, these results confirmed alterations in the immunophenotype of DC differentiated in the presence of FOH. Even though altered expression of CD1a, CD40 and HLA-DR induced by FOH could not be restored by treatment with different NR antagonists, the altered expression of CD1d, CD80 and CD86 observed after differentiation was partially dependent on PPAR γ and RAR α activation.




Figure 16. CD80 and CD40 molecule expression modulated by FOH in presence of nuclear receptors antagonists

Monocytes were differentiated into DC in the presence of either 50 μ M or 100 μ M FOH or mock treatment and the nuclear receptor antagonists indicated. For each surface marker, median surface expression after 6 days of differentiation (iDC) and 24 hours stimulation with LPS (mDC) was quantified by flow cytometry. Quantitative data shown represent means \pm SD from at least 5 independent experiments with cells isolated from different donors (* P < 0.05; ** P < 0.01; *** P < 0.001). PPAR γ antagonist: GW9662, RAR α antagonist: AGN193109, LXR α antagonist: GSK2033.




Figure 17. CD86 and HLA-DR molecule expression modulated by FOH in presence of nuclear receptors antagonists

Monocytes were differentiated into DC in the presence of either 50 µM or 100 µM FOH or mock treatment and the nuclear receptor antagonists indicated. For each surface marker, median surface expression after 6 days of differentiation (iDC) and 24 hours stimulation with LPS (mDC) was quantified by flow cytometry. Quantitative data shown represent means \pm SD from at least 5 independent experiments with cells isolated from different donors (* P < 0.05; ** P < 0.01; *** P < 0.001). PPAR γ antagonist: GW9662, RAR α antagonist: AGN193109, LXR α antagonist: GSK2033.

6.3.1.4 Release of cytokines and chemokines by farnesol-differentiated DC is mainly independent of nuclear receptors activation

The release of cytokines and chemokines by DC play an important role in regulating T cell activation. Cells were differentiated in presence or absence of NR antagonists and then stimulated with LPS to test the contribution of NR activity in the secretion of cytokines and chemokines (Figure 18). IL-8 and MCP-1 (CCL2) secretion was enhanced in FOH-differentiated DC and not influenced by NR antagonists. Interestingly, the FOH-induced increased secretion of pro-inflammatory TNF- α was blocked in presence of LXR α antagonist (100 μ M FOH: without LXR α antagonist, 8.7 ± 3.5 ng/ml; with LXR α antagonist, 3.0 ± 1.4 ng/ml, $P < 0.05$) and restored TNF α levels to those observed in mock-treated DC (4.4 ± 1.0 ng/ml). In contrast, cells treated with RAR α antagonist showed a reduction in TNF- α in every tested condition, suggesting a general and FOH-independent effect. Interestingly, differentiation of DC with either PPAR γ , RAR α or LXR α antagonists in the absence of FOH resulted in elevated levels of the anti-inflammatory cytokine IL-10, and synergistically increased its release in cells differentiated with 50 μ M FOH. However, IL-10 release induced by higher concentration of FOH was not further enhanced by blocked NR activities. In contrast, the release of RANTES was dampened in FOH-instructed DC in a concentration-dependent manner and the presence of PPAR γ antagonist partially increases the release of this chemokine in every tested condition. Moreover, IL-12 secretion was impaired in FOH-differentiated DC and treatment with the respective antagonists was not sufficient to restore the secretion of this cytokine by mDC. Treatment with the LXR α antagonist alone inhibited IL-12 secretion in mock-treated mDC and during exposure to the lower FOH concentration.




Figure 18. Cytokines and chemokines release by DC differentiated in presence of FOH and NR antagonists

Supernatants were collected after stimulation of iDC that were differentiated either in the absence or presence of FOH (50 μM and 100 μM) for 24 hours with LPS. Nuclear receptor activities were blocked using the respective antagonists: PPARγ antagonist GW9662; RARα antagonist AGN193109; LXRα antagonist GSK2033. Cytokine and chemokines concentrations were determined by multiplex assay. Data are means ± SD of 4 independent experiments using cells isolated from different donors (* P < 0.05; ** P < 0.01; *** P < 0.001).

6.3.2 Role of MAPK and NF- κ B signaling cascades in farnesol-differentiated DC

Pathway analysis of differentially expressed genes revealed that numerous processes were influenced by FOH stimulation, including antigen processing and presentation, PPAR signaling pathways, cytokine-cytokine interaction, and NF- κ B signaling (56). In particular, activation of MAPK and NF- κ B signaling pathways is known to be essential for proper differentiation and maturation of DC from monocytes (74, 75, 78, 79). To obtain further insights of the molecular mechanisms that regulate the altered maturation of DC, cell surface markers and cytokines release were measured from iDC treated with specific NF- κ B and MAPK inhibitors prior stimulation with LPS.

6.3.2.1 MAPK and NF- κ B signaling pathways modulate maturation of DC

Treatment of DC with MAPK and NF- κ B inhibitors did not restore CD1a downregulation in FOH-differentiated DC (Figure 19). Interestingly, upregulation of CD1d by FOH was partially blocked in presence of p38 inhibitor (mDC with 100 μ M FOH, median of 473 ± 84 ; mDC with 100 μ M FOH and p38 inhibitor, median of 341 ± 52 , p : 0.07). As expected, the upregulation of CD80 by LPS stimulation was significantly blocked in the presence of p38 MAPK and NF- κ B inhibitors (mock-treated mDC, median of 1102 ± 165 ; with NF- κ B inhibitor, median of 778 ± 171 , $P < 0.01$; with p38 inhibitor, median of 645 ± 100 , $P < 0.001$). Moreover, during FOH treatment we observed a synergistic reduction on the expression of CD80 in the presence of p38 MAPK and NF- κ B inhibitors (mDC with 50 μ M FOH, median of 810 ± 168 ; mDC with 50 μ M FOH and NF- κ B inhibitor, median of 559 ± 150 , $P < 0.05$; mDC with 50 μ M FOH and p38 inhibitor, median of 409 ± 86 , $P < 0.001$). In contrast, CD40 and CD86 surface exposure in DC-differentiated in absence or presence of FOH was blocked only in the presence of the NF- κ B inhibitor. Finally, the surface exposure of HLA-DR was not affected in any of the conditions tested.




Figure 19. Surface molecule expression in FOH-differentiated mDC in presence of MAPK and NF-κB inhibitors

Monocytes were differentiated into iDC in the presence of either 50 μM or 100 μM FOH or mock treatment followed by stimulation with LPS for 24 hours. For each surface marker, median surface was quantified by flow cytometry. Quantitative data represent means ± SD from 3 independent experiments with cells isolated from different donors (* $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$). MAPK and NF-κB activation were blocked using respective antagonists: SC75741: NF-κB inhibitor, SP600125: JNK inhibitor, SB203580: p38 inhibitor, FR180204: ERK1/2 inhibitor.

6.3.2.2 Farnesol regulates cytokine and chemokine secretion through MAPK and NF- κ B signaling pathways

Our previous results showed that FOH modulate the expression CD1d and costimulatory molecules through activation of NF- κ B and p38 MAPK. Thereby, we proceeded our analysis by measuring the release of cytokines and chemokines in DC differentiated in presence or absence of FOH, NF- κ B and MAPK inhibitors. Even though the release of IL-8 was not blocked by any of the inhibitors tested (Figure 20), the presence of NF- κ B inhibitor blocked the enhanced release of MCP-1, TNF- α and IL-10 induced by FOH. Interestingly, treatment with JNK and p38 MAPK inhibitors impaired the elevated secretion of TNF- α and IL-10 induced by FOH, respectively. Moreover, reduction in RANTES secretion was observed in presence of NF- κ B, JNK and p38 inhibitors in every tested condition. Similarly, NF- κ B and p38 signaling pathways also regulated IL-12 release in mock-treated mDC (mock-treated mDC, 3217 ± 1379 pg/ml; mock-treated mDC with NF- κ B inhibitor, 1206 ± 796 pg/ml, $P < 0.05$; mock-treated mDC with p38 inhibitor, 61 ± 24 pg/ml, $P < 0.001$). In contrast, during FOH treatment a synergistic reduction was observed in the presence of the p38 inhibitor (50 μ M FOH, 402 ± 221 pg/ml; 50 μ M FOH and p38 inhibitor, 27 ± 13 pg/ml). To confirm if FOH treatment modify the activation of these signaling pathways, phosphorylation levels of p65-NF- κ B and p38 MAPK was evaluated through flow cytometry. PhosphoFlow analysis showed that iDC differentiated in the presence of FOH have a higher activation of NF- κ B compared to mock-treated DC (0 min with 100 μ M FOH, fold change of 3 ± 0.3 ; without 100 μ M FOH, fold change of 2.5 ± 0.4 , $P < 0.05$) (Figure 21). As expected, phosphorylation of p65 increased after 60 min of LPS stimulation in mock-treated DC, but no differences were observed compared to FOH-differentiated DC. Finally, DC differentiated in presence of 100 μ M FOH showed an enhanced p38 phosphorylation compared to mock-treated DC before and after 60 min of LPS stimulation (Figure 22). Altogether, these results indicate that FOH modulates activation of MAPK and NF- κ B signaling pathways which results in altered expression of costimulatory molecules and cytokine release.




Figure 20. Cytokine and chemokine release by DC differentiated in presence of FOH and NF-κB and MAPK pathway inhibitors

iDC were differentiated either in the absence or presence of FOH (50 μM and 100 μM), stimulated with LPS for 24 hours, and then supernatants were collected. MAPK and NF-κB activation were blocked using respective antagonists: SC75741: NF-κB inhibitor, SP600125: JNK inhibitor, SB203580: p38 inhibitor, FR180204: ERK1/2 inhibitor. Cytokine concentrations were determined by multiplex assay. Data are means ± SD of 3 independent experiments using cells isolated from different donors (* $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$).




Figure 21. Analysis of NF- κ B activation in FOH-differentiated mDC

iDC were differentiated either in the absence (mock-treated) or presence of FOH (100 μ M) and phosphorylation of p65 NF- κ B was addressed through flow cytometry. Figure shows representative histogram of p38 phosphorylation followed by statistical analysis of DC before (A) and after 60 min of LPS stimulation (B). Quantitative analysis is from 4 independent experiments with cells isolated from different donors and shown as fold change values acquired between DC-differentiated in absence or presence of FOH and normalized to values observed in unstained DC (* $P < 0.05$).




Figure 22. Analysis of p38 activation in FOH-differentiated mDC

iDC were differentiated either in the absence (mock-treated) or presence of FOH (100 µM) and phosphorylation of p38 MAPK was addressed through flow cytometry. Figure shows representative histogram of p38 phosphorylation followed by statistical analysis of DC before (A) and after 60 min of LPS stimulation (B). Quantitative analysis is from 4 independent experiments with cells isolated from different donors and shown as fold change values acquired between DC-differentiated in absence or presence of FOH and normalized to values observed in unstained DC (* $P < 0.05$).

6.4 Functional impact of farnesol treatment in DC

DC play an important role in orchestrating immune response through antigen-presentation and activation of effector T cells (65). Consequently, alteration of these functions might have important consequences in the ability to overcome infection by pathogenic microorganisms. To evaluate if FOH treatment alters DC capacity to activate T cells, mDC and T cell co-cultures were performed and the ability of mDC to activate iNKT, Th1 and Treg was addressed by flow cytometry.

6.4.1 Farnesol-differentiated DC have a lower capacity to activate iNKT and Th1 cells

The previous results showed that FOH alters dramatically the expression of CD1d molecules through PPAR γ , RAR α and partially through p38 MAPK activation. CD1d is a major-histocompatibility-complex (MHC) class-1-like molecule that presents lipid antigens to invariant Natural Killer T (iNKT) cells to activate them (85, 87). To test whether the elevated CD1d surface levels induced by FOH would correlate with an increased capability of mDC to induce iNKT activation, expansion of iNKT cells was analyzed by flow cytometry. Surprisingly, iNKT proliferation was reduced in the presence of FOH-differentiated mDC (50 μ M FOH, fold change of 2.5 ± 1.2 , $P < 0.01$) compared to the mock-treated mDC (Fold change of 7.3 ± 1.7), regardless of their significantly enhanced CD1d levels (Figure 23). Although CD1d expression induced by FOH was dependent on both PPAR γ and RAR α activity, blocking NR activities resulted in comparably low iNKT expansion.

All data reported so far suggest that FOH-differentiated mDC have a defect to induce proper T cell responses, by reducing the expression of costimulatory molecules and the MHC class II molecule HLA-DR. To evaluate if FOH alters the capacity of mDC to activate Th1 cells, mDC differentiated in presence or absence of FOH and NR antagonists were co-cultured with autologous T cells for 7 days and IFN- γ production was analyzed by flow cytometry after restimulation with PMA/ionomycin. As expected, T cells co-cultured with mock-treated mDC (Fold change of 16.6 ± 5.4) have a higher ability to induce IFN- γ production compared to FOH differentiated mDC (50 μ M FOH, fold change of 5.2 ± 2.7 , $P < 0.05$) (Figure 24). The presence of specific antagonists for PPAR γ and RAR α did not restore the capability of mDC to induce IFN- γ production by T cells.




Figure 23. iNKT expansion by FOH treated-DC

Autologous T cells were co-cultured with mDC for 7 days. iNKT expansion was measured by flow cytometry and quantitative analysis from 4 independent experiments with cells isolated from different donors is displayed. Quantitative analysis is shown as means \pm SD of fold change values acquired between % iNKT cells of total T cells in co-culture with mock- or FOH-treated mDC normalized to values observed in T cells cultured without mDC (* P < 0.05; ** P < 0.01). GW: PPAR γ antagonist GW9662; AGN: RAR α antagonist AGN193109.




Figure 24. IFN- γ production by T cells co-cultured with mDC

Autologous T cells were co-cultured with mDC for 7 days. T cells were collected and re-stimulated with PMA (50 ng/mL) and ionomycin (1 μ g/mL), in the presence of brefeldin A (5 μ g/mL) and monensin (5 μ g/mL) for 6 hours. IFN- γ production was analyzed by intracellular staining and flow cytometry. Quantitative analysis of 3 independent experiments is shown as means \pm SD of fold change values acquired between % IFN- γ ⁺ cells of total T cells co-culture with mock- or FOH-treated mDC normalized to values observed in T cells cultured without mDC (* P < 0.05; ** P < 0.01). PPAR γ antagonist: GW9662; RAR α antagonist: AGN193109.

6.4.1.1 Altered secretion of IL-12 and IL-10 influences the capacity of farnesol-differentiated DC to activate different T cell subsets

Although FOH increased CD1d expression in DC, no positive effect was observed in their capacity to induce iNKT expansion. A possible explanation could be the increased IL-10 secretion and reduced IL-12 release observed in FOH-treated mDC. Due to their potent effect on activating T cells, IL-12 family members are an important link between innate and adaptive immunity. Moreover, IL-10 is known to inhibit IL-12 production by macrophages and DC (206). To test if the shift in the IL-12/IL-10 ratio observed in FOH-differentiated DC might play a role in iNKT activation (Figure 25A), T cell-mDC co-culture experiments were performed in the presence of increased IL-12 levels by addition of recombinant IL-12, while IL-10 activity was reduced by the use of blocking antibody. Interestingly, the reduced iNKT proliferation observed in presence of FOH-differentiated DC was increased after reconstitution of the IL-12/IL-10 milieu by addition of recombinant IL-12 and neutralizing α IL-10, reaching levels similar to the mock T cell-mDC co-culture (mock-treated mDC, fold change of 10.1 ± 3.7 , with 50 μ M FOH, fold change of 4.7 ± 2.9 ; 50 μ M FOH and IL-12/ α IL-10, fold change of 12.1 ± 4.8 , $P < 0.05$) (Figure 25B). Furthermore, addition of recombinant IL-12 and α IL-10 also enhanced iNKT proliferation in mock T cell-mDC co-culture (Fold change of 15.4 ± 3.8).

To test if the altered IL-12 and IL-10 release also played a role in mDC capability to induce IFN- γ production by T cells, co-culture of mDC and autologous T cells for 7 days was performed and IFN- γ production by T cells was induced by restimulation with PMA/ionomycin (Figure 26). In line with the result observed for iNKT expansion, the reduced percentage of IFN- γ producing T cells co-cultured with FOH-differentiated mDC was restored to levels observed in mock-treated mDC co-culture (Fold change of 6.8 ± 4.9) when elevated concentrations of IL-12 and IL-10 blocking antibody were present (50 μ M FOH, fold change of 1.3 ± 0.3 ; 50 μ M FOH and IL-12/ α IL-10, fold change of 6.7 ± 4.8 , $P < 0.05$).



A**B**

Figure 25. Diminished capacity to induce iNKT expansion by FOH-differentiated mDC is dependent on the altered secretion of IL-12 and IL-10

iDC were differentiated either in the absence or presence of FOH (50 µM and 100 µM), stimulated with LPS for 24 hours, and then supernatants were collected. (A) Data show the ratio between the measured concentrations of IL-12 and IL-10. (B) iNKT expansion was analyzed by flow cytometry after 7 days of co-culture with mock-treated or FOH-instructed mDC in the presence of recombinant IL-12 and anti-IL-10 mAb or the respective isotype control. Bars are means \pm SD from 4 independent experiments using cells isolated from different donors and show fold change values acquired between % iNKT cells of total T cells in co-culture with mock- or FOH-treated mDC normalized to values observed in T cells cultured without mDC (* P < 0.05; ** P < 0.01, *** P < 0.001).




Figure 26. Altered release of IL-12 and IL-10 by FOH-treated mDC impacts their capacity to induce T helper 1 polarization

Autologous T cells were co-cultured with mock-treated or FOH-differentiated mDC in presence or absence of IL-12 and anti-IL-10 mAb, or an isotype control, and IFN γ production by T cells was evaluated after 7 days of co-culture. Bars are means \pm SD from 4 independent experiments using cells isolated from different donors and show fold change values between % IFN γ + T cells of total T cells in co-culture with mock- or FOH-treated mDC and normalized to values observed in T cells cultured without mDC (*P < 0.05).

6.4.2 Farnesol-differentiated DC have a higher expression of inhibitory receptors but lack capacity to induce regulatory T cells expansion

The low expression of costimulatory molecules, together with a higher secretion of IL-10 and reduced capacity of FOH-differentiated mDC to activate iNKT and Th1 cells, might indicate that FOH treatment shift DC functionality towards a tolerogenic phenotype. Therefore, analysis of inhibitory receptors expression, including Ig-like transcripts (ILTs, also called leukocyte Ig-like receptors) ILT2, ILT3, ILT4, and programmed cell death 1 ligand 1 (PD-L1), known to be a hallmark of tolerogenic DC (107, 109), was performed by flow cytometry after differentiation and 24 hours LPS stimulation. The expression of ILT2 and ILT4 was not influenced by FOH (Figure 27). Nevertheless, increased surface exposure of ILT3 after LPS stimulation was evidenced in FOH-differentiated mDC compared to mock-treated cells (mDC with 100 μ M FOH, median of 12000 ± 1032 , mock-treated mDC, median of 3716 ± 549 ; $P < 0.001$). Similarly, expression of PD-L1 was enhanced in mDC differentiated in presence of FOH (mDC with 100 μ M FOH, median of 4740 ± 151 ; mock-treated mDC, median of 3136 ± 482 ; $P < 0.001$). Elevated expression of inhibitory receptors and IL-10 secretion was hypothesized to enhance the capacity to activate FOXP3⁺ Treg cells (104, 207). Thereby, T cells were co-cultured with mDC differentiated in absence or presence of FOH and Treg cells expansion was evaluated after 7 days. Interestingly, the enhanced expression of inhibitory receptors in FOH-differentiated mDC did not lead to elevated Treg cells expansion compared to mock-treated mDC (mDC with 100 μ M FOH, fold change 1.8 ± 1.5 ; mock-treated mDC, fold change 13.6 ± 10.5) (Figure 28). In line with our previous results, the addition of recombinant IL-12 and IL-10 blocking antibody restored FOH-differentiated DC capacity to activate Treg cells to comparable levels observed in mock-treated mDC co-culture (mDC with 100 μ M FOH and IL-12/ α IL-10, fold change 11.6 ± 6.5).




Figure 27. FOH enhances the surface expression of inhibitory receptors on mDC

Monocytes were differentiated into iDC in the presence or absence of FOH followed by a 24 hours LPS stimulation to generate mDC. The expression of ILT2, ILT3, ILT4 and PD-L1 was analyzed by flow cytometry. Quantitative data are means \pm SD from at least 4 independent experiments with cells isolated from different donors (***) $P < 0.001$ compared to mock-treated).




Figure 28. FOH altered DC capacity to induce FOXP3+ regulatory T cells expansion

Autologous T cells were co-cultured with mDC for 7 days in the absence or presence of recombinant IL-12 and anti-IL-10 mAb, or the respective isotype control and CD4⁺CD25⁺FOXP3⁺ Treg cells expansion was measured by flow cytometry. Quantitative data are means \pm SD from 4 independent experiments with cells isolated from different donors ($***P < 0.001$).

7 Discussion

QS systems coordinate the expression of genes involved in several processes, including virulence in pathogenic microorganisms (15, 208). FOH is the first QS molecule described in a eukaryotic microorganism, and its role in modulating filamentation in *C. albicans* is well understood (27-32). Similarly, FOH has been shown to regulate functionality of innate immune cells, such as neutrophils, monocytes, macrophages and dendritic cells. Leonhardt *et al.* showed that FOH has a profound impact on dendritic cells differentiation from monocytes, while suppressing their capacity to induce T cell proliferation. However, the molecular basis behind these effects is poorly understood (56).

The aim of this study was to analyze the signaling pathways modulated by FOH during dendritic cells differentiation, and its impact on their function.

7.1 Farnesol impairs DC differentiation and maturation partially through nuclear receptors activation

Leonhardt *et al.* found that FOH restrict the differentiation from monocytes to DC, by retaining the expression of the monocyte marker CD14 while impairing costimulatory molecules surface exposure and T cell activation capacity (56). This study confirmed that FOH blocked maturation of DC by modifying the expression of costimulatory and antigen-presenting molecules, as well as the release of cytokines and chemokines in response to *C. albicans* germ tubes and several ligands to Toll-like receptors and dectin-1. These results suggest that FOH-induced changes are not dependent of any specific surface receptor and affect intracellular signaling cascades. Pathway analysis of differentially expressed genes during DC differentiation identified several intracellular processes that were modified in presence of FOH, including the PPAR signaling pathway (56). In agreement, transcriptional analysis revealed that several target genes associated to activation of NR PPAR γ , RAR α and LXR α were upregulated. Moreover, differentiation of DC in presence of specific ligands for these NR showed a similar phenotype compared to FOH-treated DC. Furthermore, the presence of PPAR γ and RAR α antagonists blocked upregulation of CD1d by FOH-treatment, and partially reverted FOH effect on CD80 and CD86 surface molecules expression and the release of the chemokine RANTES. Interestingly, RAR α and LXR α

showed to have a major role in the release of pro-inflammatory TNF- α induced by FOH. While these results suggest that FOH activates NR, it is not clear how these cells sense FOH in the environment. On the one hand, FOH could be interacting directly with NR as suggested by nuclear localization of a fluorescent analog (20). On the other hand, FOH can be metabolized into farnesyl-PP, an intermediate of the mevalonate/cholesterol biosynthetic pathway, which can be further transformed into NR ligands (209). However, Shchepin *et al.* showed that the mode of action of FOH is specific, since changes in its structure blocked its biological activity (20). Interestingly, FOH is not the only QS molecule known to affect DC differentiation. Treatment of DC with the bacterial QS molecule homoserine lactone diminished the expression of the costimulatory molecules CD80 and CD86 (210). Interestingly, the effect of QS molecules on DC differentiation could have an important impact on septic patients. Boontham *et al.* showed that homoserine lactone effects on DC function correlated with elevated presence of this QS molecule in sera of septic patients (211). Although it is not known the role of FOH during sepsis, Ryan *et al.* showed that monocyte-derived DC from autoimmune polyendocrinopathy-candidiasis-ectodermal dystrophy (APECED) had impaired DC maturation (212). Similar to our results, these DC showed enhanced production of TNF- α and IL-8 and reduced capacity to prime T cells (56, 212).

7.2 Farnesol regulation of CD1 molecules and its functional impact

The most prominent effect observed after dendritic cells differentiation in presence of FOH was the displacement in the expression of CD1 molecules. While CD1a expression was not upregulated, CD1d was enhanced during DC differentiation in the presence of FOH (56). CD1 molecules present lipid antigens to different T cells subsets. In particular, CD1a are specialized in presenting lipid antigens to T cells, while CD1d present antigens to iNKT cells (85). iNKT cells are a subpopulation of T cells with the capacity to modulate immune responses due to its capability to produce high amounts of pro-inflammatory and anti-inflammatory cytokines. Thereby, the shift in the CD1 molecules expression pattern may influence the outcome of infection, since iNKT cells are known to be important during infection against pathogenic microorganisms (213-215). However, there are discrepancies regarding the role of iNKT cells during *C. albicans* infection. Cohen *et al.* showed that iNKT cells activation reduces disease progression in a murine model of systemic candidiasis

(213). Similarly, iNKT cells recognize *C. albicans* α -mannosyl residues in a CD1d-dependent manner and increases the survival of mice co-infected with *S. pneumoniae* (215). Nevertheless, different reports showed that iNKT cells have no role or that activation of these cells promotes susceptibility to systemic *C. albicans* infection due to high IL-10 production (216, 217). Thus, more research to elucidate the role of iNKT cells during *C. albicans* infection is needed. While a higher expression of CD1d correlates with an elevated capacity to activate iNKT cells, we failed to observe an enhanced expansion of these cells when co-cultured with FOH-differentiated mDC (87, 156). One possibility could be that FOH-differentiated DC have an impaired capacity to present antigens to T cells. Nakken *et al.* showed that endosomal processing of antigens by proteases is necessary for proper induction of iNKT expansion by DC (178). However, the antigen used in this study (α -GalCer) does not require preprocessing and readily binds to CD1d complex (218). Thereby, a different mechanism should be involved in the reduced capacity of FOH-mDC to activate iNKT cells. In concert with signals triggered by CD1d-mediated lipid presentation, costimulatory molecule interactions between DC and T cells, such as CD40:CD40L and CD80/CD86:CD28, and the release of IL-12 by DC is required for optimal iNKT expansion (219, 220). Indeed, a reduction in CD40 and CD80 was observed in DC differentiated with FOH. However, our results indicate that the shift in the IL-12/IL-10 secretion ratio play an essential role in FOH-differentiated mDC capacity to induce iNKT expansion, since reconstitution of IL-12 and IL-10 to levels observed in absence of FOH significantly improves DC capability to expand iNKT cells.

This work gives insights into the molecular mechanisms by which FOH modulates CD1d expression in DC. Our results showed that FOH regulates CD1d expression via activation of the PPAR α /RAR α signaling pathway. Szatmari *et al.* demonstrated that activation of PPAR γ leads to upregulation of retinaldehyde dehydrogenases (RALDH, ALDH1) that catalyze the production of all-trans retinoic acid (ATRA) (156). ATRA serves as a ligand for RAR α which increases CD1d expression in human DC (156). However, proper lipid antigen processing and loading to CD1d molecules requires activation of the lysosomal aspartyl protease cathepsin D CTSD (178). Concurrent with the higher CD1d surface exposure on FOH-differentiated DC, analysis of the transcriptional data identified increased gene expression of NR target genes, including ALDH1A1 and CTSD. Our results confirm previous observations that FOH activates NR, such as PPAR γ (209, 221-223). Mechanistically, FOH

could interact directly or indirectly with NR (209, 224). Interestingly, our data suggest that FOH could regulate NR activities through activation of MAPK signaling pathway. Indeed, inhibition of p38 MAPK partially restored CD1d basal levels in FOH-differentiated mDC. NR activities can be modulated independently of the presence of ligands through activation of the AF-1 N-terminal region, which serves as target for post-translational modifications, including phosphorylation (127, 128, 130, 131). Moreover, p38 inhibition showed to diminished PPAR γ transcriptional activities in different cell types (225, 226). Finally, we found that FOH-differentiated DC have a higher activation of p38 MAPK compared to mock-treated DC. Altogether, these results indicate that FOH modulates CD1d surface expression via activation of the PPAR γ /RAR α signaling pathway in a process partially dependent on p38 MAPK.

7.3 Farnesol alters cytokines release through several pathways

The results presented in this study confirm that FOH modulates cytokines and chemokines release by mDC. While production of pro-inflammatory cytokines/chemokines (e.g. TNF- α , IL-8, MCP-1), and the anti-inflammatory cytokine IL-10 was enhanced in FOH-differentiated mDC, T cell activating cytokine and chemokine (e.g. IL-12 and RANTES) was reduced after LPS stimulation. We found that FOH modulates the release of cytokines through a complex activation of NR, MAPK and NF- κ B signaling pathways. In particular, TNF- α production in FOH-differentiated mDC was blocked in the presence of inhibitors for JNK MAPK, NF- κ B, RAR α and LXR α . Accordingly, Szanto *et al.* showed that RAR α activation by natural and synthetic ligands promotes the expression of genes such as CYP27A1, an enzyme involved in ligand production for LXR α (227). Activation of LXR α also activates NF- κ B, which enhances the secretion of pro-inflammatory TNF- α (191). Interestingly, the elevated release of anti-inflammatory IL-10 observed in FOH-differentiated mDC might be a possible mechanism to suppress the inflammatory milieu observed in our experiments. Indeed, IL-10 production was blocked when p38 MAPK and NF- κ B inhibitors were present in the culture and correlated with the secretion of pro-inflammatory cytokines and chemokines observed in FOH-differentiated mDC. Moreover, analyses of p38 MAPK and NF- κ B phosphorylation confirmed that FOH-instructed mDC have a higher activation of these pathways compared to mock-treated mDC. However, the diminished secretion of IL-12 and RANTES indicate that FOH-instructed mDC have a reduced T cell activation capacity. Although RANTES

production was partially dependent on PPAR γ activation, our results suggest that FOH suppresses IL-12 release through a MAPK, NF- κ B and NR independent mechanism. IL-12 gene expression is regulated by MAPK/AP-1, NF- κ B and IRF signaling pathways (228). Previous results showed that IL-12p70 release was reduced in IRF-1 deficient DC (228, 229). Moreover, TLR4 activation leads to IRF-3 recruitment to the IL-12p35 promoter, which enhances IL-12p70 synthesis (230). Thus, it is possible that FOH impairs IL-12p70 production in mDC by modulating IRF activation. Interestingly, other microbial QS molecules are known to affect IL-12 release. Homoserine lactone and the *Pseudomona aeruginosa* quinolone signal have been shown to decrease IL-12 production by murine DC (210). Moreover, another molecule produced by *C. albicans*, CA-SIIF, is known to reduce CD1a expression and IL-12 secretion in human DC (231). These results suggest that IL-12 reduction might be a mechanism used by *C. albicans* to overcome immune responses orchestrated by DC.

In summary, our results showed that FOH promotes inflammation, while reduces T cell inducing cytokines and chemokines, in a process dependent on several signaling pathways, including NR, MAPK and NF- κ B.

7.4 Farnesol-treated DC have a lower capacity to activate different T cell subsets

DC are considered to be the bridge between the innate and adaptive branch of immunity, due to its potent capacity to present antigens and consequently activating T cell-specific responses (65). The altered expression of costimulatory molecules and cytokines release induced by FOH might indicate an impaired capacity of FOH-differentiated DC to activate T cells. Indeed, FOH-instructed mDC have a lower capability to activate Th1, iNKT and Treg cells. IFN- γ produced by Th1 lymphocytes is essential for stimulation of antifungal activity by neutrophils (97). The role of IFN- γ in anti-*Candida* immunity is highlighted by the enhanced mortality observed in IFN- γ -deficient mice infected with *C. albicans* and the elevated susceptibility of HIV-infected individuals with low CD4⁺ T cell count to develop oropharyngeal candidiasis (232, 233). Moreover, IFN- γ contributes to antifungal immunity by inducing nitric oxide production by macrophages, while reducing *Candida*-specific immunoglobulins (234). In agreement with our data, the bacteria QS molecule homoserine

lactone and *P. aeruginosa* quinolone signal has been shown to alter DC capacity to induce activation of T cells (210, 211). However, this is the first report that shows a negative effect of QS molecules in DC capacity to activate FOXP3⁺ Treg cells. While Th1 cells orchestrate antifungal immunity mainly by activating phagocytes, Treg are involved in controlling inflammation by suppressing the secretion of pro-inflammatory cytokines (235). Indeed, Treg expand during disseminated infection and produce high amounts of IL-4, TGF- β and IL-10, which blocks Th1 immunity (236, 237). Surprisingly, Treg could improve immunity against pathogenic fungi via an alternative mechanism. Treg was shown to promote Th17 differentiation during systemic candidemia and oropharyngeal candidiasis (98, 236). IL-17 play an important role in anti-fungal immunity and exerts its function mainly through recruitment and activation of neutrophils (98, 238). The reduced expression of costimulatory molecules observed in FOH-instructed mDC might influence the impaired capability of these cells to activate T cells. However, our results showed that surface molecules play a minor role in DC capability to induce Treg activation, since the enhanced expression of inhibitory receptors, such as ILT3 and PD-L1 did not correlate with increased FOXP3⁺ Treg expansion. Reconstitution of IL-12 and IL-10 release to normal levels was sufficient to restore the capacity of FOH-differentiated mDC to activate Th1 and Treg populations. IL-12 is an important cytokine involved in the differentiation and activation of T cells. IL-12 signals through the IL-12 receptor, triggers activation of the Janus kinase 2 (JAK2) and tyrosine kinase 2 (TYK2), leading to phosphorylation of family members of the STAT transcription factor, resulting in polarization of T cells into INF- γ producing Th1 cells (239). Interestingly, IL-12 signaling increased IFN- γ production in Treg while retaining FOXP3⁺ expression (240).

Altogether, our results confirm that FOH acts as a compound with immune modulatory properties that blocks the capacity of DC to induce protective adaptive immunity. Since T cells play an important role during *C. albicans* infection, our results support the role of FOH as a virulence factor produce by this fungus to evade DC surveillance.

8 Conclusions

Besides its well-known quorum sensing properties, FOH has been shown to be a modulator of immune function. FOH is able to activate monocytes, neutrophils and macrophages. Interestingly, this molecule altered DC differentiation and maturation while reducing their capacity to activate T cells. Due to the relevance of DC in orchestrating immune responses during infection, identification of the molecular mechanisms by which FOH regulates DC function could lead to possible targets for pharmacological strategies against *C. albicans* infection. Our findings showed for the first time that FOH modulates DC differentiation and maturation through activation of the nuclear receptors PPAR γ , RAR α and LXR α , MAPK and NF- κ B signaling pathways (Figure 29). FOH activates the PPAR γ /RAR α axis to increase CD1d surface expression on DC in a process partially dependent on p38 MAPK signaling. We found that FOH increased the release of pro-inflammatory cytokines and anti-inflammatory IL-10, and impaired IL-12 release mainly through the RAR α , LXR α , MAPK and NF- κ B signaling pathways. Despite multiple effects of FOH that in isolation suggest activation of DC, the net effect of FOH exposure is paralysis of DC function due to the reduced ability to prime several T cell subsets. In agreement, the altered secretion of IL-12 and IL-10 induced by FOH treatment showed to play a key role in FOH-differentiated DC capacity to promote iNKT, Th1 and Treg cell activation. Due to the relevance of these cells in the anti-Candida immunity, besides its quorum sensing function, FOH can be viewed as a virulence factor enabling *C. albicans* to overcome the immune surveillance by DC.

Further studies should focus in elucidating the molecular mechanisms underlying the diminished production of IL-12 in FOH-differentiated DC. One possibility might be to evaluate the IRF signaling pathway which is known to modulate IL-12 release in DC (228-230). Furthermore, it would be interesting to characterize the interaction with *Candida albicans* in respect to phagocytosis and killing capacity. The current work shows evidences of the crosstalk between host nuclear receptors and molecules produced by *C. albicans*. Interestingly, fungi present nuclear-receptor like transcription factors (241, 242); thus, it would be interesting to elucidate the role of these transcription factors in the virulence of *C. albicans* and evaluate how molecules produced by host cells might interact with these transcription factors.




Figure 29. Farnesol modulates DC function through a coordinated activation of several signaling pathways

FOH activates several intracellular signaling pathways resulting in a distinct DC subset, with altered expression of antigen-presenting and costimulatory molecules. FOH modulates the expression of CD1d via p38 MAPK-PPAR γ /RAR α signaling pathway. Moreover, FOH-differentiated DC showed an elevated secretion of pro-inflammatory and anti-inflammatory IL-10, through MAPK and NF- κ B activation. Finally, FOH treatment reduces DC capacity to activate different T cell subsets, such as Th1, iNKT and Treg cells, via reduction of the IL-12/IL-10 secretion ratio.

9 References

1. Mayer, F.L., D. Wilson, and B. Hube, *Candida albicans pathogenicity mechanisms*. Virulence, 2013. **4**(2): p. 119-28.
2. Noble, S.M., B.A. Gianetti, and J.N. Witchley, *Candida albicans cell-type switching and functional plasticity in the mammalian host*. Nature Reviews Microbiology, 2016. **15**: p. 96.
3. Perlroth, J., B. Choi, and B. Spellberg, *Nosocomial fungal infections: epidemiology, diagnosis, and treatment*. Med Mycol, 2007. **45**(4): p. 321-46.
4. Moyes, D.L., et al., *Candidalysin is a fungal peptide toxin critical for mucosal infection*. Nature, 2016. **532**(7597): p. 64-8.
5. Berman, J. and P.E. Sudbery, *Candida Albicans: a molecular revolution built on lessons from budding yeast*. Nat Rev Genet, 2002. **3**(12): p. 918-30.
6. Sudbery, P.E., *Growth of Candida albicans hyphae*. Nat Rev Microbiol, 2011. **9**(10): p. 737-48.
7. Garcia, M.C., et al., *A Role for Amyloid in Cell Aggregation and Biofilm Formation*. PLOS ONE, 2011. **6**(3): p. e17632.
8. Nobile, C.J. and A.D. Johnson, *Candida albicans Biofilms and Human Disease*. Annu Rev Microbiol, 2015. **69**: p. 71-92.
9. Phan, Q.T., et al., *Als3 is a Candida albicans invasin that binds to cadherins and induces endocytosis by host cells*. PLoS Biol, 2007. **5**(3): p. e64.
10. Phan, Q.T., et al., *N-cadherin mediates endocytosis of Candida albicans by endothelial cells*. J Biol Chem, 2005. **280**(11): p. 10455-61.
11. Chen, H., et al., *Tyrosol is a quorum-sensing molecule in Candida albicans*. Proc Natl Acad Sci U S A, 2004. **101**(14): p. 5048-52.
12. Oh, K.B., et al., *Purification and characterization of an autoregulatory substance capable of regulating the morphological transition in Candida albicans*. Proc Natl Acad Sci U S A, 2001. **98**(8): p. 4664-8.
13. Hornby, J.M., et al., *Quorum sensing in the dimorphic fungus Candida albicans is mediated by farnesol*. Applied and environmental microbiology, 2001. **67**(7): p. 2982-2992.
14. Alem, M.A., et al., *Production of tyrosol by Candida albicans biofilms and its role in quorum sensing and biofilm development*. Eukaryot Cell, 2006. **5**(10): p. 1770-9.
15. Polke, M., et al., *Farnesol signalling in Candida albicans - more than just communication*. Crit Rev Microbiol, 2018. **44**(2): p. 230-243.
16. Lv, Q.-Z., L. Yan, and Y.-Y. Jiang, *The synthesis, regulation, and functions of sterols in Candida albicans: Well-known but still lots to learn*. Virulence, 2016. **7**(6): p. 649-659.
17. Nickerson, K.W., A.L. Atkin, and J.M. Hornby, *Quorum sensing in dimorphic fungi: farnesol and beyond*. Applied and environmental microbiology, 2006. **72**(6): p. 3805-3813.
18. Hornby, J.M. and K.W. Nickerson, *Enhanced Production of Farnesol by Candida albicans Treated with Four Azoles*. Antimicrobial Agents and Chemotherapy, 2004. **48**(6): p. 2305.
19. Hornby, J.M., B.W. Kebaara, and K.W. Nickerson, *Farnesol Biosynthesis in Candida albicans: Cellular Response to Sterol Inhibition by Zaragozic Acid B*. Antimicrobial Agents and Chemotherapy, 2003. **47**(7): p. 2366.
20. Shchepin, R., et al., *Quorum sensing in Candida albicans: probing farnesol's mode of action with 40 natural and synthetic farnesol analogs*. Chem Biol, 2003. **10**(8): p. 743-50.

21. Weber, K., et al., *Secretion of E,E-farnesol and biofilm formation in eight different Candida species*. Antimicrobial agents and chemotherapy, 2008. **52**(5): p. 1859-1861.
22. Navarathna, D.H., et al., *Effect of farnesol on a mouse model of systemic candidiasis, determined by use of a DPP3 knockout mutant of Candida albicans*. Infect Immun, 2007. **75**(4): p. 1609-18.
23. Zhang, L., et al., *Bisbibenzyls, a New Type of Antifungal Agent, Inhibit Morphogenesis Switch and Biofilm Formation through Upregulation of DPP3 in Candida albicans*. PLOS ONE, 2011. **6**(12): p. e28953.
24. Kadosh, D. and A.D. Johnson, *Induction of the Candida albicans filamentous growth program by relief of transcriptional repression: a genome-wide analysis*. Mol Biol Cell, 2005. **16**(6): p. 2903-12.
25. Kebaara, B.W., et al., *Candida albicans Tup1 is involved in farnesol-mediated inhibition of filamentous-growth induction*. Eukaryot Cell, 2008. **7**(6): p. 980-7.
26. Polke, M., et al., *A functional link between hyphal maintenance and quorum sensing in Candida albicans*. Mol Microbiol, 2017. **103**(4): p. 595-617.
27. Hornby, J.M., et al., *Quorum sensing in the dimorphic fungus Candida albicans is mediated by farnesol*. Appl Environ Microbiol, 2001. **67**(7): p. 2982-92.
28. Lindsay, A.K., et al., *Farnesol and cyclic AMP signaling effects on the hypha-to-yeast transition in Candida albicans*. Eukaryot Cell, 2012. **11**(10): p. 1219-25.
29. Ramage, G., et al., *Inhibition of Candida albicans biofilm formation by farnesol, a quorum-sensing molecule*. Applied and environmental microbiology, 2002. **68**(11): p. 5459-5463.
30. Uppuluri, P., et al., *Dispersion as an important step in the Candida albicans biofilm developmental cycle*. PLoS Pathog, 2010. **6**(3): p. e1000828.
31. Cao, Y.Y., et al., *cDNA microarray analysis of differential gene expression in Candida albicans biofilm exposed to farnesol*. Antimicrob Agents Chemother, 2005. **49**(2): p. 584-9.
32. Cho, T., et al., *Transcriptional changes in Candida albicans Genes by both farnesol and high cell density at an early stage of morphogenesis in N-acetyl-D-glucosamine medium*. Nihon Ishinkin Gakkai Zasshi, 2007. **48**(4): p. 159-67.
33. Feng, Q., et al., *Ras signaling is required for serum-induced hyphal differentiation in Candida albicans*. Journal of bacteriology, 1999. **181**(20): p. 6339-6346.
34. Piispanen, A.E., et al., *Regulated proteolysis of Candida albicans Ras1 is involved in morphogenesis and quorum sensing regulation*. Mol Microbiol, 2013. **89**(1): p. 166-78.
35. Kuroda, Y., N. Suzuki, and T. Kataoka, *The effect of posttranslational modifications on the interaction of Ras2 with adenylyl cyclase*. Science, 1993. **259**(5095): p. 683-6.
36. Rocha, C.R., et al., *Signaling through adenylyl cyclase is essential for hyphal growth and virulence in the pathogenic fungus Candida albicans*. Mol Biol Cell, 2001. **12**(11): p. 3631-43.
37. Langford, M.L., A.L. Atkin, and K.W. Nickerson, *Cellular interactions of farnesol, a quorum-sensing molecule produced by Candida albicans*. Future Microbiol, 2009. **4**(10): p. 1353-62.
38. Hall, R.A., et al., *The quorum-sensing molecules farnesol/homoserine lactone and dodecanol operate via distinct modes of action in Candida albicans*. Eukaryot Cell, 2011. **10**(8): p. 1034-42.
39. Roman, E., et al., *The Cek1 MAPK is a short-lived protein regulated by quorum sensing in the fungal pathogen Candida albicans*. FEMS Yeast Res, 2009. **9**(6): p. 942-55.
40. Kruppa, M., et al., *The two-component signal transduction protein Chk1p regulates quorum sensing in Candida albicans*. Eukaryot Cell, 2004. **3**(4): p. 1062-5.

41. Langford, M.L., et al., *Candida albicans Czf1 and Efg1 coordinate the response to farnesol during quorum sensing, white-opaque thermal dimorphism, and cell death*. Eukaryotic cell, 2013. **12**(9): p. 1281-1292.
42. Lorek, J., et al., *Influence of farnesol on the morphogenesis of Aspergillus niger*. J Basic Microbiol, 2008. **48**(2): p. 99-103.
43. Semighini, C.P., N. Murray, and S.D. Harris, *Inhibition of Fusarium graminearum growth and development by farnesol*. FEMS Microbiol Lett, 2008. **279**(2): p. 259-64.
44. Liu, P., et al., *Farnesol induces apoptosis and oxidative stress in the fungal pathogen Penicillium expansum*. Mycologia, 2010. **102**(2): p. 311-8.
45. Fairn, G.D., K. Macdonald, and C.R. McMaster, *A chemogenomic screen in Saccharomyces cerevisiae uncovers a primary role for the mitochondria in farnesol toxicity and its regulation by the Pkc1 pathway*. J Biol Chem, 2007. **282**(7): p. 4868-74.
46. Wang, X., et al., *Farnesol induces apoptosis-like cell death in the pathogenic fungus Aspergillus flavus*. Mycologia, 2014. **106**(5): p. 881-888.
47. Semighini, C.P., et al., *Farnesol-induced apoptosis in Aspergillus nidulans reveals a possible mechanism for antagonistic interactions between fungi*. Mol Microbiol, 2006. **59**(3): p. 753-64.
48. Dichtl, K., et al., *Farnesol misplaces tip-localized Rho proteins and inhibits cell wall integrity signalling in Aspergillus fumigatus*. Mol Microbiol, 2010. **76**(5): p. 1191-204.
49. Brehm-Stecher, B.F. and E.A. Johnson, *Sensitization of Staphylococcus aureus and Escherichia coli to antibiotics by the sesquiterpenoids nerolidol, farnesol, bisabolol, and apritone*. Antimicrob Agents Chemother, 2003. **47**(10): p. 3357-60.
50. Jabra-Rizk, M.A., et al., *Effect of farnesol on Staphylococcus aureus biofilm formation and antimicrobial susceptibility*. Antimicrobial agents and chemotherapy, 2006. **50**(4): p. 1463-1469.
51. Scheper, M.A., et al., *Farnesol, a fungal quorum-sensing molecule triggers apoptosis in human oral squamous carcinoma cells*. Neoplasia, 2008. **10**(9): p. 954-63.
52. Joo, J.H., et al., *Farnesol-induced apoptosis in human lung carcinoma cells is coupled to the endoplasmic reticulum stress response*. Cancer Res, 2007. **67**(16): p. 7929-36.
53. Park, J.S., et al., *Farnesol induces apoptosis of DU145 prostate cancer cells through the PI3K/Akt and MAPK pathways*. Int J Mol Med, 2014. **33**(5): p. 1169-76.
54. Duncan, R.E. and M.C. Archer, *Farnesol induces thyroid hormone receptor (THR) beta1 but inhibits THR-mediated signaling in MCF-7 human breast cancer cells*. Biochem Biophys Res Commun, 2006. **343**(1): p. 239-43.
55. Rennemeier, C., et al., *Microbial quorum-sensing molecules induce acrosome loss and cell death in human spermatozoa*. Infection and immunity, 2009. **77**(11): p. 4990-4997.
56. Leonhardt, I., et al., *The fungal quorum-sensing molecule farnesol activates innate immune cells but suppresses cellular adaptive immunity*. MBio, 2015. **6**(2): p. e00143.
57. Abe, S., et al., *Suppression of anti-Candida activity of macrophages by a quorum-sensing molecule, farnesol, through induction of oxidative stress*. 2009. **53**(6): p. 323-330.
58. Navarathna, D.H., et al., *Enhanced pathogenicity of Candida albicans pre-treated with subinhibitory concentrations of fluconazole in a mouse model of disseminated candidiasis*. J Antimicrob Chemother, 2005. **56**(6): p. 1156-9.
59. Tashiro, M., et al., *Pravastatin inhibits farnesol production in Candida albicans and improves survival in a mouse model of systemic candidiasis*. Med Mycol, 2012. **50**(4): p. 353-60.

60. Hargarten, J.C., et al., *Candida albicans Quorum Sensing Molecules Stimulate Mouse Macrophage Migration*. Infection and Immunity, 2015. **83**(10): p. 3857.
61. Ghosh, S., et al., *Candida albicans cell wall components and farnesol stimulate the expression of both inflammatory and regulatory cytokines in the murine RAW264.7 macrophage cell line*. FEMS Immunol Med Microbiol, 2010. **60**(1): p. 63-73.
62. Hunniger, K. and O. Kurzai, *Phagocytes as central players in the defence against invasive fungal infection*. Semin Cell Dev Biol, 2019. **89**: p. 3-15.
63. Duggan, S., et al., *Host response to Candida albicans bloodstream infection and sepsis*. Virulence, 2015. **6**(4): p. 316-26.
64. Collin, M. and V. Bigley, *Human dendritic cell subsets: an update*. Immunology, 2018. **154**(1): p. 3-20.
65. Thery, C. and S. Amigorena, *The cell biology of antigen presentation in dendritic cells*. Curr Opin Immunol, 2001. **13**(1): p. 45-51.
66. Swiecki, M. and M. Colonna, *The multifaceted biology of plasmacytoid dendritic cells*. Nat Rev Immunol, 2015. **15**(8): p. 471-85.
67. Embgenbroich, M. and S. Burgdorf, *Current Concepts of Antigen Cross-Presentation*. Frontiers in immunology, 2018. **9**: p. 1643-1643.
68. Banchereau, J. and R.M. Steinman, *Dendritic cells and the control of immunity*. Nature, 1998. **392**(6673): p. 245-52.
69. Kimber, I. and M. Cumberbatch, *Stimulation of Langerhans cell migration by tumor necrosis factor alpha (TNF-alpha)*. J Invest Dermatol, 1992. **99**(5): p. 48s-50s.
70. De Smedt, T., et al., *Regulation of dendritic cell numbers and maturation by lipopolysaccharide in vivo*. J Exp Med, 1996. **184**(4): p. 1413-24.
71. Guo, J. and S.L. Friedman, *Toll-like receptor 4 signaling in liver injury and hepatic fibrogenesis*. Fibrogenesis & Tissue Repair, 2010. **3**(1): p. 21.
72. Lu, Y.C., W.C. Yeh, and P.S. Ohashi, *LPS/TLR4 signal transduction pathway*. Cytokine, 2008. **42**(2): p. 145-151.
73. Kawai, T. and S. Akira, *TLR signaling*. Cell Death & Differentiation, 2006. **13**(5): p. 816-825.
74. Rescigno, M., et al., *Dendritic Cell Survival and Maturation Are Regulated by Different Signaling Pathways*. J Exp Med, 1998. **188**(11): p. 2175-2180.
75. Hernandez, A., et al., *Inhibition of NF-kappa B during human dendritic cell differentiation generates anergy and regulatory T-cell activity for one but not two human leukocyte antigen DR mismatches*. Hum immunol, 2007. **68**(9): p. 715-729.
76. van de Laar, L., et al., *A nonredundant role for canonical NF-kappaB in human myeloid dendritic cell development and function*. J Immunol, 2010. **185**(12): p. 7252-7261.
77. Andreaskos, E., et al., *Activation of NF-kappaB by the intracellular expression of NF-kappaB-inducing kinase acts as a powerful vaccine adjuvant*. Proc Natl Acad Sci U S A, 2006. **103**(39): p. 14459-14464.
78. Arrighi, J.F., et al., *A critical role for p38 mitogen-activated protein kinase in the maturation of human blood-derived dendritic cells induced by lipopolysaccharide, TNF-alpha, and contact sensitizers*. J Immunol, 2001. **166**(6): p. 3837-3845.
79. Nakahara, T., et al., *Role of c-Jun N-terminal kinase on lipopolysaccharide induced maturation of human monocyte-derived dendritic cells*. Int Immunol, 2004. **16**(12): p. 1701-1709.

80. Kasinrerker, W., et al., *CD1 molecule expression on human monocytes induced by granulocyte-macrophage colony-stimulating factor*. J Immunol, 1993. **150**(2): p. 579-84.
81. Sallusto, F. and A. Lanzavecchia, *Efficient presentation of soluble antigen by cultured human dendritic cells is maintained by granulocyte/macrophage colony-stimulating factor plus interleukin 4 and downregulated by tumor necrosis factor alpha*. J Exp Med, 1994. **179**(4): p. 1109-18.
82. Romani, N., et al., *Generation of mature dendritic cells from human blood. An improved method with special regard to clinical applicability*. J Immunol Methods, 1996. **196**(2): p. 137-51.
83. Hopkins, R.A. and J.E. Connolly, *The specialized roles of immature and mature dendritic cells in antigen cross-presentation*. Immunologic Research, 2012. **53**(1): p. 91-107.
84. Van Kaer, L., L. Wu, and S. Joyce, *Mechanisms and Consequences of Antigen Presentation by CD1*. Trends Immunol, 2016. **37**(11): p. 738-754.
85. Brigl, M. and M.B. Brenner, *CD1: antigen presentation and T cell function*. Annu Rev Immunol, 2004. **22**: p. 817-90.
86. Angenieux, C., et al., *Characterization of CD1e, a third type of CD1 molecule expressed in dendritic cells*. J Biol Chem, 2000. **275**(48): p. 37757-64.
87. Szatmari, I., et al., *Activation of PPARgamma specifies a dendritic cell subtype capable of enhanced induction of iNKT cell expansion*. Immunity, 2004. **21**(1): p. 95-106.
88. Carreño, L.J., Shalu S. Kharkwal, and S.A. Porcelli, *Optimizing NKT cell ligands as vaccine adjuvants*. 2014. **6**(3): p. 309-320.
89. Hubo, M., et al., *Costimulatory Molecules on Immunogenic Versus Tolerogenic Human Dendritic Cells*. 2013. **4**(82).
90. Lenschow, D.J., T.L. Walunas, and J.A. Bluestone, *CD28/B7 system of T cell costimulation*. Annu Rev Immunol, 1996. **14**: p. 233-58.
91. Thiel, M., et al., *Efficiency of T-cell costimulation by CD80 and CD86 cross-linking correlates with calcium entry*. Immunology, 2010. **129**(1): p. 28-40.
92. Linsley, P.S., et al., *Binding of the B cell activation antigen B7 to CD28 costimulates T cell proliferation and interleukin 2 mRNA accumulation*. J Exp Med, 1991. **173**(3): p. 721-30.
93. Grewal, I.S. and R.A. Flavell, *CD40 and CD154 in cell-mediated immunity*. Annu Rev Immunol, 1998. **16**: p. 111-35.
94. Luckheeram, R.V., et al., *CD4⁺T cells: differentiation and functions*. Clinical & developmental immunology, 2012. **2012**: p. 925135-925135.
95. Trinchieri, G., S. Pflanz, and R.A. Kastelein, *The IL-12 family of heterodimeric cytokines: new players in the regulation of T cell responses*. Immunity, 2003. **19**(5): p. 641-4.
96. Walker, J.A. and A.N.J. McKenzie, *TH2 cell development and function*. Nature Reviews Immunology, 2017. **18**: p. 121.
97. van de Veerdonk, F.L. and M.G. Netea, *T-cell Subsets and Antifungal Host Defenses*. Current fungal infection reports, 2010. **4**(4): p. 238-243.
98. Pandiyan, P., et al., *CD4(+)CD25(+)Foxp3(+) regulatory T cells promote Th17 cells in vitro and enhance host resistance in mouse Candida albicans Th17 cell infection model*. Immunity, 2011. **34**(3): p. 422-34.
99. Robinson, M.J., et al., *Dectin-2 is a Syk-coupled pattern recognition receptor crucial for Th17 responses to fungal infection*. J Exp Med, 2009. **206**(9): p. 2037-51.
100. Hernández-Santos, N. and S.L. Gaffen, *Th17 cells in immunity to Candida albicans*. Cell host & microbe, 2012. **11**(5): p. 425-435.

101. Vignali, D.A.A., L.W. Collison, and C.J. Workman, *How regulatory T cells work*. Nature Reviews Immunology, 2008. **8**: p. 523.
102. Hori, S., T. Nomura, and S. Sakaguchi, *Control of regulatory T cell development by the transcription factor Foxp3*. Science, 2003. **299**(5609): p. 1057-61.
103. Fontenot, J.D., M.A. Gavin, and A.Y. Rudensky, *Foxp3 programs the development and function of CD4+CD25+ regulatory T cells*. Nat Immunol, 2003. **4**(4): p. 330-6.
104. Maldonado, R.A. and U.H. von Andrian, *How tolerogenic dendritic cells induce regulatory T cells*. Adv Immunol, 2010. **108**: p. 111-65.
105. Takenaka, M.C. and F.J. Quintana, *Tolerogenic dendritic cells*. Seminars in Immunopathology, 2017. **39**(2): p. 113-120.
106. Mascanfroni, I.D., et al., *IL-27 acts on DCs to suppress the T cell response and autoimmunity by inducing expression of the immunoregulatory molecule CD39*. Nat Immunol, 2013. **14**(10): p. 1054-63.
107. Sim, W.J., P.J. Ahl, and J.E. Connolly, *Metabolism Is Central to Tolerogenic Dendritic Cell Function*. Mediators Inflamm, 2016. **2016**: p. 2636701.
108. Riley, J.L., *PD-1 signaling in primary T cells*. Immunological reviews, 2009. **229**(1): p. 114-125.
109. Wu, J. and A. Horuzsko, *Expression and function of immunoglobulin-like transcripts on tolerogenic dendritic cells*. Human immunology, 2009. **70**(5): p. 353-356.
110. Qin, Y., et al., *Innate immune cell response upon Candida albicans infection*. Virulence, 2016. **7**(5): p. 512-526.
111. Kashem, S.W., et al., *Candida albicans morphology and dendritic cell subsets determine T helper cell differentiation*. Immunity, 2015. **42**(2): p. 356-366.
112. Newman, S.L. and A. Holly, *Candida albicans Is Phagocytosed, Killed, and Processed for Antigen Presentation by Human Dendritic Cells*. Infection and Immunity, 2001. **69**(11): p. 6813.
113. d'Ostiani, C.F., et al., *Dendritic cells discriminate between yeasts and hyphae of the fungus Candida albicans. Implications for initiation of T helper cell immunity in vitro and in vivo*. The Journal of experimental medicine, 2000. **191**(10): p. 1661-1674.
114. van de Veerdonk, F.L., et al., *Th17 responses and host defense against microorganisms: an overview*. BMB Rep, 2009. **42**(12): p. 776-87.
115. Whitney, P.G., et al., *Syk signaling in dendritic cells orchestrates innate resistance to systemic fungal infection*. PLoS Pathog, 2014. **10**(7): p. e1004276.
116. Pietrella, D., et al., *Candida albicans mannoprotein influences the biological function of dendritic cells*. Cell Microbiol, 2006. **8**(4): p. 602-12.
117. Netea, M.G., et al., *Role of TLR1 and TLR6 in the host defense against disseminated candidiasis*. FEMS Immunol Med Microbiol, 2008. **52**(1): p. 118-23.
118. del Fresno, C., et al., *Interferon-beta production via Dectin-1-Syk-IRF5 signaling in dendritic cells is crucial for immunity to C. albicans*. Immunity, 2013. **38**(6): p. 1176-86.
119. Cambi, A., et al., *The C-type lectin DC-SIGN (CD209) is an antigen-uptake receptor for Candida albicans on dendritic cells*. Eur J Immunol, 2003. **33**(2): p. 532-8.
120. Fermin Lee, A., et al., *Galectin-3 modulates Th17 responses by regulating dendritic cell cytokines*. Am J Pathol, 2013. **183**(4): p. 1209-1222.
121. Cambi, A., et al., *Dendritic cell interaction with Candida albicans critically depends on N-linked mannan*. The Journal of biological chemistry, 2008. **283**(29): p. 20590-20599.

122. Forsyth, C.B., E.F. Plow, and L. Zhang, *Interaction of the fungal pathogen Candida albicans with integrin CD11b/CD18: recognition by the I domain is modulated by the lectin-like domain and the CD18 subunit*. J Immunol, 1998. **161**(11): p. 6198-205.
123. Romani, L., *Immunity to fungal infections*. Nature Reviews Immunology, 2011. **11**: p. 275.
124. Saijo, S. and Y. Iwakura, *Dectin-1 and Dectin-2 in innate immunity against fungi*. Int Immunol, 2011. **23**(8): p. 467-72.
125. Tang, J., et al., *Regulation of C-Type Lectin Receptor-Mediated Antifungal Immunity*. 2018. **9**(123).
126. Plato, A., J.A. Willment, and G.D. Brown, *C-type lectin-like receptors of the dectin-1 cluster: ligands and signaling pathways*. Int Rev Immunol, 2013. **32**(2): p. 134-56.
127. Nagy, L., et al., *Nuclear hormone receptors enable macrophages and dendritic cells to sense their lipid environment and shape their immune response*. Physiol Rev, 2012. **92**(2): p. 739-89.
128. Kiss, M., Z. Czimmerer, and L. Nagy, *The role of lipid-activated nuclear receptors in shaping macrophage and dendritic cell function: From physiology to pathology*. J Allergy Clin Immunol, 2013. **132**(2): p. 264-86.
129. Huang, W. and C.K. Glass, *Nuclear receptors and inflammation control: molecular mechanisms and pathophysiological relevance*. Arterioscler Thromb Vasc Biol, 2010. **30**(8): p. 1542-9.
130. Mazaira, G.I., et al., *The Nuclear Receptor Field: A Historical Overview and Future Challenges*. Nuclear receptor research, 2018. **5**: p. 101320.
131. Rastinejad, F., et al., *Understanding nuclear receptor form and function using structural biology*. J Mol Endocrinol, 2013. **51**(3): p. T1-t21.
132. Germain, P., et al., *Overview of nomenclature of nuclear receptors*. Pharmacol Rev, 2006. **58**(4): p. 685-704.
133. Germain, P., et al., *International Union of Pharmacology. LXIII. Retinoid X Receptors*. 2006. **58**(4): p. 760-772.
134. Leid, M., et al., *Purification, cloning, and RXR identity of the HeLa cell factor with which RAR or TR heterodimerizes to bind target sequences efficiently*. Cell, 1992. **68**(2): p. 377-95.
135. Heyman, R.A., et al., *9-cis retinoic acid is a high affinity ligand for the retinoid X receptor*. Cell, 1992. **68**(2): p. 397-406.
136. Dawson, M.I. and Z. Xia, *The retinoid X receptors and their ligands*. Biochimica et biophysica acta, 2012. **1821**(1): p. 21-56.
137. Szeles, L., et al., *Research resource: transcriptome profiling of genes regulated by RXR and its permissive and nonpermissive partners in differentiating monocyte-derived dendritic cells*. Mol Endocrinol, 2010. **24**(11): p. 2218-31.
138. Zapata-Gonzalez, F., et al., *9-cis-Retinoic acid (9cRA), a retinoid X receptor (RXR) ligand, exerts immunosuppressive effects on dendritic cells by RXR-dependent activation: inhibition of peroxisome proliferator-activated receptor gamma blocks some of the 9cRA activities, and precludes them to mature phenotype development*. J Immunol, 2007. **178**(10): p. 6130-9.
139. Villablanca, E.J., et al., *Selected natural and synthetic retinoids impair CCR7- and CXCR4-dependent cell migration in vitro and in vivo*. J Leukoc Biol, 2008. **84**(3): p. 871-9.
140. Huang, Z., et al., *Role of Vitamin A in the Immune System*. Journal of clinical medicine, 2018. **7**(9): p. 258.

141. Aibana, O., et al., *Impact of Vitamin A and Carotenoids on the Risk of Tuberculosis Progression*. Clin Infect Dis, 2017. **65**(6): p. 900-909.
142. Shivakoti, R., et al., *Prevalence and risk factors of micronutrient deficiencies pre- and post-antiretroviral therapy (ART) among a diverse multicountry cohort of HIV-infected adults*. Clin Nutr, 2016. **35**(1): p. 183-9.
143. Qi, Y.J., et al., *Relationship between deficiencies in vitamin A and E and occurrence of infectious diseases among children*. Eur Rev Med Pharmacol Sci, 2016. **20**(23): p. 5009-5012.
144. So, E.N. and D.L. Crowe, *Characterization of a retinoic acid responsive element in the human ets-1 promoter*. IUBMB Life, 2000. **50**(6): p. 365-70.
145. Mendelsohn, C., et al., *Function of the retinoic acid receptors (RARs) during development (II). Multiple abnormalities at various stages of organogenesis in RAR double mutants*. Development, 1994. **120**(10): p. 2749-71.
146. Kastner, P. and S. Chan, *Function of RARalpha during the maturation of neutrophils*. Oncogene, 2001. **20**(49): p. 7178-85.
147. Sucov, H.M. and R.M.J.M.N. Evans, *Retinoic acid and retinoic acid receptors in development*. 1995. **10**(2): p. 169-184.
148. Vellozo, N.S., et al., *All-Trans Retinoic Acid Promotes an M1- to M2-Phenotype Shift and Inhibits Macrophage-Mediated Immunity to Leishmania major*. Front Immunol, 2017. **8**: p. 1560.
149. Wang, X., C. Allen, and M. Ballow, *Retinoic acid enhances the production of IL-10 while reducing the synthesis of IL-12 and TNF-alpha from LPS-stimulated monocytes/macrophages*. J Clin Immunol, 2007. **27**(2): p. 193-200.
150. Mehta, K., et al., *Inhibition by all-trans-retinoic acid of tumor necrosis factor and nitric oxide production by peritoneal macrophages*. J Leukoc Biol, 1994. **55**(3): p. 336-42.
151. Tao, Y., Y. Yang, and W. Wang, *Effect of all-trans-retinoic acid on the differentiation, maturation and functions of dendritic cells derived from cord blood monocytes*. FEMS Immunol Med Microbiol, 2006. **47**(3): p. 444-50.
152. Saurer, L., K.C. McCullough, and A. Summerfield, *In vitro induction of mucosa-type dendritic cells by all-trans retinoic acid*. J Immunol, 2007. **179**(6): p. 3504-14.
153. Bakdash, G., et al., *Retinoic acid primes human dendritic cells to induce gut-homing, IL-10-producing regulatory T cells*. Mucosal Immunology, 2014. **8**: p. 265.
154. Manicassamy, S., et al., *Toll-like receptor 2-dependent induction of vitamin A-metabolizing enzymes in dendritic cells promotes T regulatory responses and inhibits autoimmunity*. Nat Med, 2009. **15**(4): p. 401-9.
155. Geissmann, F., et al., *Retinoids regulate survival and antigen presentation by immature dendritic cells*. The Journal of experimental medicine, 2003. **198**(4): p. 623-634.
156. Szatmari, I., et al., *PPARgamma controls CD1d expression by turning on retinoic acid synthesis in developing human dendritic cells*. J Exp Med, 2006. **203**(10): p. 2351-62.
157. Darmanin, S., et al., *All-trans retinoic acid enhances murine dendritic cell migration to draining lymph nodes via the balance of matrix metalloproteinases and their inhibitors*. J Immunol, 2007. **179**(7): p. 4616-25.
158. Klassert, T.E., et al., *Modulatory role of vitamin A on the Candida albicans-induced immune response in human monocytes*. Medical microbiology and immunology, 2014. **203**(6): p. 415-424.

159. Tyagi, S., et al., *The peroxisome proliferator-activated receptor: A family of nuclear receptors role in various diseases*. Journal of advanced pharmaceutical technology & research, 2011. **2**(4): p. 236-240.
160. Grygiel-Górniak, B.J.N.J., *Peroxisome proliferator-activated receptors and their ligands: nutritional and clinical implications - a review*. 2014. **13**(1): p. 17.
161. Pawlak, M., P. Lefebvre, and B. Staels, *Molecular mechanism of PPAR α action and its impact on lipid metabolism, inflammation and fibrosis in non-alcoholic fatty liver disease*. Journal of Hepatology, 2015. **62**(3): p. 720-733.
162. Chinetti, G., et al., *Activation of proliferator-activated receptors alpha and gamma induces apoptosis of human monocyte-derived macrophages*. J Biol Chem, 1998. **273**(40): p. 25573-80.
163. Jones, D.C., X. Ding, and R.A. Daynes, *Nuclear receptor peroxisome proliferator-activated receptor alpha (PPARalpha) is expressed in resting murine lymphocytes. The PPARalpha in T and B lymphocytes is both transactivation and transrepression competent*. J Biol Chem, 2002. **277**(9): p. 6838-45.
164. Palomer, X., et al., *PPAR δ : A Key Therapeutic Target in Metabolic Disorders*. International journal of molecular sciences, 2018. **19**(3): p. 913.
165. Széles, L., D. Töröcsik, and L. Nagy, *PPAR γ in immunity and inflammation: cell types and diseases*. Biochimica et Biophysica Acta (BBA) - Molecular and Cell Biology of Lipids, 2007. **1771**(8): p. 1014-1030.
166. Nolte, R.T., et al., *Ligand binding and co-activator assembly of the peroxisome proliferator-activated receptor-gamma*. Nature, 1998. **395**(6698): p. 137-43.
167. Jiang, C., A.T. Ting, and B. Seed, *PPAR-gamma agonists inhibit production of monocyte inflammatory cytokines*. Nature, 1998. **391**(6662): p. 82-6.
168. Alleva, D.G., et al., *Regulation of murine macrophage proinflammatory and anti-inflammatory cytokines by ligands for peroxisome proliferator-activated receptor-gamma: counter-regulatory activity by IFN-gamma*. J Leukoc Biol, 2002. **71**(4): p. 677-85.
169. Chung, S.W., et al., *Oxidized low density lipoprotein inhibits interleukin-12 production in lipopolysaccharide-activated mouse macrophages via direct interactions between peroxisome proliferator-activated receptor-gamma and nuclear factor-kappa B*. J Biol Chem, 2000. **275**(42): p. 32681-7.
170. Welch, J.S., et al., *PPARgamma and PPARdelta negatively regulate specific subsets of lipopolysaccharide and IFN-gamma target genes in macrophages*. Proc Natl Acad Sci U S A, 2003. **100**(11): p. 6712-7.
171. Tanaka, T., et al., *Therapeutic potential of thiazolidinediones in activation of peroxisome proliferator-activated receptor gamma for monocyte recruitment and endothelial regeneration*. Eur J Pharmacol, 2005. **508**(1-3): p. 255-65.
172. Ricote, M. and C.K. Glass, *PPARs and molecular mechanisms of transrepression*. Biochimica et biophysica acta, 2007. **1771**(8): p. 926-935.
173. Szanto, A., et al., *STAT6 transcription factor is a facilitator of the nuclear receptor PPARgamma-regulated gene expression in macrophages and dendritic cells*. Immunity, 2010. **33**(5): p. 699-712.
174. Appel, S., et al., *PPAR- γ agonists inhibit toll-like receptor-mediated activation of dendritic cells via the MAP kinase and NF- κ B pathways*. 2005. **106**(12): p. 3888-3894.

175. Zhu, W., et al., *PPAR-gamma agonist pioglitazone regulates dendritic cells immunogenicity mediated by DC-SIGN via the MAPK and NF-kappaB pathways*. *Int Immunopharmacol*, 2016. **41**: p. 24-34.
176. Gosset, P., et al., *Peroxisome proliferator-activated receptor gamma activators affect the maturation of human monocyte-derived dendritic cells*. *Eur J Immunol*, 2001. **31**(10): p. 2857-65.
177. Nencioni, A., et al., *Dendritic cell immunogenicity is regulated by peroxisome proliferator-activated receptor gamma*. *J Immunol*, 2002. **169**(3): p. 1228-35.
178. Nakken, B., et al., *Peroxisome proliferator-activated receptor gamma-regulated cathepsin D is required for lipid antigen presentation by dendritic cells*. *J Immunol*, 2011. **187**(1): p. 240-7.
179. Guirado, E., et al., *Deletion of PPARgamma in lung macrophages provides an immunoprotective response against M. tuberculosis infection in mice*. *Tuberculosis (Edinb)*, 2018. **111**: p. 170-177.
180. Gautier, E.L., et al., *Systemic analysis of PPARgamma in mouse macrophage populations reveals marked diversity in expression with critical roles in resolution of inflammation and airway immunity*. *J Immunol*, 2012. **189**(5): p. 2614-24.
181. Bedi, B., et al., *Enhanced Clearance of Pseudomonas aeruginosa by Peroxisome Proliferator-Activated Receptor Gamma*. *Infect Immun*, 2016. **84**(7): p. 1975-1985.
182. Majer, O., et al., *Type I interferons promote fatal immunopathology by regulating inflammatory monocytes and neutrophils during Candida infections*. *PLoS Pathog*, 2012. **8**(7): p. e1002811.
183. Gales, A., et al., *PPARgamma controls Dectin-1 expression required for host antifungal defense against Candida albicans*. *PLoS Pathog*, 2010. **6**(1): p. e1000714.
184. Lefevre, L., et al., *PPARgamma ligands switched high fat diet-induced macrophage M2b polarization toward M2a thereby improving intestinal Candida elimination*. *PLoS One*, 2010. **5**(9): p. e12828.
185. Coste, A., et al., *IL-13 attenuates gastrointestinal candidiasis in normal and immunodeficient RAG-2(-/-) mice via peroxisome proliferator-activated receptor-gamma activation*. *J Immunol*, 2008. **180**(7): p. 4939-47.
186. Waddington, Kirsty E., Elizabeth C. Jury, and I. Pineda-Torra, *Liver X receptors in immune cell function in humans*. 2015. **43**(4): p. 752-757.
187. Schulman, I.G., *Liver X receptors link lipid metabolism and inflammation*. 2017. **591**(19): p. 2978-2991.
188. Joseph, S.B., et al., *LXR-dependent gene expression is important for macrophage survival and the innate immune response*. *Cell*, 2004. **119**(2): p. 299-309.
189. Pourcet, B., et al., *The nuclear receptor LXR modulates interleukin-18 levels in macrophages through multiple mechanisms*. *Scientific Reports*, 2016. **6**: p. 25481.
190. Geyeregger, R., et al., *Liver X receptors regulate dendritic cell phenotype and function through blocked induction of the actin-bundling protein fascin*. *Blood*, 2007. **109**(10): p. 4288-95.
191. Torocsik, D., et al., *Activation of liver X receptor sensitizes human dendritic cells to inflammatory stimuli*. *J Immunol*, 2010. **184**(10): p. 5456-5465.
192. Beceiro, S., et al., *Liver X Receptor Nuclear Receptors Are Transcriptional Regulators of Dendritic Cell Chemotaxis*. *Mol Cell Biol*, 2018. **38**(10).

193. Castrillo, A., et al., *Liver X receptor-dependent repression of matrix metalloproteinase-9 expression in macrophages*. J Biol Chem, 2003. **278**(12): p. 10443-9.
194. Joseph, S.B., et al., *Reciprocal regulation of inflammation and lipid metabolism by liver X receptors*. Nat Med, 2003. **9**(2): p. 213-9.
195. Ghisletti, S., et al., *Parallel SUMOylation-dependent pathways mediate gene- and signal-specific transrepression by LXRs and PPARgamma*. Molecular cell, 2007. **25**(1): p. 57-70.
196. Fontaine, C., et al., *Liver X receptor activation potentiates the lipopolysaccharide response in human macrophages*. Circ Res, 2007. **101**(1): p. 40-9.
197. Frasca, L., et al., *CD38 orchestrates migration, survival, and Th1 immune response of human mature dendritic cells*. Blood, 2006. **107**(6): p. 2392-9.
198. Korf, H., et al., *Liver X receptors contribute to the protective immune response against Mycobacterium tuberculosis in mice*. J Clin Invest, 2009. **119**(6): p. 1626-37.
199. Bouttier, M., et al., *Alu repeats as transcriptional regulatory platforms in macrophage responses to M. tuberculosis infection*. Nucleic Acids Res, 2016. **44**(22): p. 10571-10587.
200. Lange, P.T., et al., *Liver X Receptors Suppress Activity of Cholesterol and Fatty Acid Synthesis Pathways To Oppose Gammaherpesvirus Replication*. MBio, 2018. **9**(4).
201. Ramezani, A., et al., *Stimulation of Liver X Receptor Has Potent Anti-HIV Effects in a Humanized Mouse Model of HIV Infection*. J Pharmacol Exp Ther, 2015. **354**(3): p. 376-83.
202. Bruhn, K.W., et al., *LXR deficiency confers increased protection against visceral Leishmania infection in mice*. PLoS Negl Trop Dis, 2010. **4**(11): p. e886.
203. Bobryshev, Y.V., et al., *Decreased expression of liver X receptor-alpha in macrophages infected with Chlamydia pneumoniae in human atherosclerotic arteries in situ*. J Innate Immun, 2011. **3**(5): p. 483-94.
204. Weber, K., B. Schulz, and M. Ruhnke, *The quorum-sensing molecule E,E-farnesol—its variable secretion and its impact on the growth and metabolism of Candida species*. Yeast, 2010. **27**(9): p. 727-739.
205. Langford, M.L., et al., *Activity and toxicity of farnesol towards Candida albicans are dependent on growth conditions*. Antimicrob Agents Ch, 2010. **54**(2): p. 940-942.
206. Ma, X., et al., *Regulation of IL-10 and IL-12 production and function in macrophages and dendritic cells*. F1000Research, 2015. **4**: p. F1000 Faculty Rev-1465.
207. Unger, W.W., et al., *Induction of Treg by monocyte-derived DC modulated by vitamin D3 or dexamethasone: differential role for PD-L1*. Eur J Immunol, 2009. **39**(11): p. 3147-59.
208. Whiteley, M., S.P. Diggle, and E.P. Greenberg, *Progress in and promise of bacterial quorum sensing research*. Nature, 2017. **551**: p. 313.
209. Joo, J.H. and A.M. Jetten, *Molecular mechanisms involved in farnesol-induced apoptosis*. Cancer Lett, 2010. **287**(2): p. 123-135.
210. Skindersoe, M.E., et al., *Pseudomonas aeruginosa quorum-sensing signal molecules interfere with dendritic cell-induced T-cell proliferation*. FEMS Immunol Med Microbiol, 2009. **55**(3): p. 335-45.
211. Boontham, P., et al., *Significant immunomodulatory effects of Pseudomonas aeruginosa quorum-sensing signal molecules: possible link in human sepsis*. Clinical Science, 2008. **115**(11): p. 343.
212. Ryan, K.R., et al., *Impaired dendritic cell maturation and cytokine production in patients with chronic mucocutaneous candidiasis with or without APECED*. Clin Exp Immunol, 2008. **154**(3): p. 406-14.

213. Cohen, N.R., et al., *Innate recognition of cell wall beta-glucans drives invariant natural killer T cell responses against fungi*. Cell Host Microbe, 2011. **10**(5): p. 437-50.
214. Fischer, K., et al., *Mycobacterial phosphatidylinositol mannoside is a natural antigen for CD1d-restricted T cells*. Proc Natl Acad Sci U S A, 2004. **101**(29): p. 10685-90.
215. Albacker, L.A., et al., *Invariant natural killer T cells recognize a fungal glycosphingolipid that can induce airway hyperreactivity*. Nat Med, 2013. **19**(10): p. 1297-304.
216. Tarumoto, N., et al., *A limited role of iNKT cells in controlling systemic Candida albicans infections*. Jpn J Infect Dis, 2012. **65**(6): p. 522-6.
217. Haraguchi, N., et al., *Activation of murine invariant NKT cells promotes susceptibility to candidiasis by IL-10 induced modulation of phagocyte antifungal activity*. Eur J Immunol, 2016. **46**(7): p. 1691-703.
218. Yu, K.O., et al., *Modulation of CD1d-restricted NKT cell responses by using N-acyl variants of alpha-galactosylceramides*. Proc Natl Acad Sci U S A, 2005. **102**(9): p. 3383-8.
219. Hayakawa, Y., et al., *Differential regulation of Th1 and Th2 functions of NKT cells by CD28 and CD40 costimulatory pathways*. J Immunol, 2001. **166**(10): p. 6012-8.
220. Kitamura, H., et al., *The natural killer T (NKT) cell ligand alpha-galactosylceramide demonstrates its immunopotentiating effect by inducing interleukin (IL)-12 production by dendritic cells and IL-12 receptor expression on NKT cells*. J Exp Med, 1999. **189**(7): p. 1121-8.
221. Forman, B.M., et al., *Identification of a nuclear receptor that is activated by farnesol metabolites*. Cell, 1995. **81**(5): p. 687-693.
222. Takahashi, N., et al., *Dual action of isoprenols from herbal medicines on both PPARgamma and PPARalpha in 3T3-L1 adipocytes and HepG2 hepatocytes*. FEBS Lett, 2002. **514**(2-3): p. 315-22.
223. Journe, F., et al., *Farnesol, a mevalonate pathway intermediate, stimulates MCF-7 breast cancer cell growth through farnesoid-X-receptor-mediated estrogen receptor activation*. Breast Cancer Res Treat, 2008. **107**(1): p. 49-61.
224. Shchepin, R., et al., *Biologically active fluorescent farnesol analogs*. Chem Biol, 2005. **12**(6): p. 639-641.
225. Schild, R.L., et al., *The kinase p38 regulates peroxisome proliferator activated receptor-gamma in human trophoblasts*. Placenta, 2006. **27**(2-3): p. 191-9.
226. Lu, Y., et al., *p38 MAPK-inhibited dendritic cells induce superior antitumour immune responses and overcome regulatory T-cell-mediated immunosuppression*. Nat Commun, 2014. **5**: p. 4229.
227. Szanto, A., et al., *Transcriptional regulation of human CYP27 integrates retinoid, peroxisome proliferator-activated receptor, and liver X receptor signaling in macrophages*. Mol Cell Biol, 2004. **24**(18): p. 8154-66.
228. Antonios, D., et al., *Mechanisms of IL-12 Synthesis by Human Dendritic Cells Treated with the Chemical Sensitizer NiSO₄*. 2010. **185**(1): p. 89-98.
229. Gabriele, L., et al., *IRF-1 deficiency skews the differentiation of dendritic cells toward plasmacytoid and tolerogenic features*. J Leukoc Biol, 2006. **80**(6): p. 1500-11.
230. Goriely, S., et al., *Interferon regulatory factor 3 is involved in Toll-like receptor 4 (TLR4)- and TLR3-induced IL-12p35 gene activation*. Blood, 2006. **107**(3): p. 1078-84.
231. Wang, M., et al., *Characterization and partial purification of Candida albicans Secretory IL-12 Inhibitory Factor*. BMC microbiology, 2008. **8**: p. 31-31.

232. Balish, E., et al., *Candidiasis in interferon-gamma knockout (IFN-gamma-/-) mice*. J Infect Dis, 1998. **178**(2): p. 478-87.
233. Vazquez, J.A., *Optimal management of oropharyngeal and esophageal candidiasis in patients living with HIV infection*. HIV/AIDS (Auckland, N.Z.), 2010. **2**: p. 89-101.
234. Kaposzta, R., et al., *Characteristics of invasive candidiasis in gamma interferon- and interleukin-4-deficient mice: role of macrophages in host defense against Candida albicans*. Infect Immun, 1998. **66**(4): p. 1708-17.
235. Vignali, D.A., L.W. Collison, and C.J. Workman, *How regulatory T cells work*. Nat Rev Immunol, 2008. **8**(7): p. 523-32.
236. Whibley, N., et al., *Expansion of Foxp3(+) T-cell populations by Candida albicans enhances both Th17-cell responses and fungal dissemination after intravenous challenge*. Eur J Immunol, 2014. **44**(4): p. 1069-83.
237. De Luca, A., et al., *Functional yet balanced reactivity to Candida albicans requires TRIF, MyD88, and IDO-dependent inhibition of Rorc*. J Immunol, 2007. **179**(9): p. 5999-6008.
238. Whibley, N. and S.L. Gaffen, *Brothers in arms: Th17 and Treg responses in Candida albicans immunity*. PLoS pathogens, 2014. **10**(12): p. e1004456-e1004456.
239. Hamza, T., J.B. Barnett, and B. Li, *Interleukin 12 a key immunoregulatory cytokine in infection applications*. Int J Mol Sci, 2010. **11**(3): p. 789-806.
240. Feng, T., et al., *Interleukin-12 converts Foxp3+ regulatory T cells to interferon-gamma-producing Foxp3+ T cells that inhibit colitis*. Gastroenterology, 2011. **140**(7): p. 2031-43.
241. Thakur, J.K., et al., *A nuclear receptor-like pathway regulating multidrug resistance in fungi*. Nature, 2008. **452**(7187): p. 604-9.
242. Naar, A.M. and J.K. Thakur, *Nuclear receptor-like transcription factors in fungi*. Genes Dev, 2009. **23**(4): p. 419-32.

10 List of abbreviations

%	Percent
°C	Degree Celsius
Ab	Antibody
AGN	RAR α antagonist
AP-1	Activator protein 1
APC	Allophycocyanin
AM	RAR α agonist
CD	Cluster of differentiation
CLR	C-type lectin receptors
DC	Dendritic cells
DMSO	Dimethylsulfoxid
DNA	Deoxyribonucleic acid
DPP3	Diacylglycerol pyrophosphate phosphatase
e.g.	exempli gratia
EDTA	Ethylenediaminetetraacetic acid
ERK	Extracellular-signal-regulated kinase
et. al.	et alii
FBS	Fetal bovine serum
FCS	Forward scatter
FITC	Fluorescein isothiocyanate
FOH	Farnesol
FOXP3	Forkhead box P3
g	Acceleration due to gravity
GM-CSF	Granulocyte macrophage colony-stimulating factor
GSK	LXR antagonist
GW	PPAR γ antagonist
HLA-DR	Human leukocyte antigen-D related
iDC	Immature dendritic cells
IFN	Interferon
IL	Interleukin
ILT	leukocyte Ig-like receptors
iNKT	Invariant Natural Killer T cells
JNK	c-Jun N-terminal kinase
LXR	Liver X receptor
LPS	Lipopolysaccharide
MAPK	Mitogen-activated protein kinase
MCP	Monocyte chemoattractant protein
mDC	Mature dendritic cells
MHC	Major histocompatibility complex
mins	Minutes
MOH	Methanol
MOI	Multiplicity of infection
NFAT	Nuclear factor activated T-cells
NF- κ B	Nuclear factor kappa B
NR	Nuclear receptor
PBMC	Peripheral blood mononuclear cell
PBS	Phosphate buffer saline
PD-L	Programmed death-ligand
PE	Phycoerythrin
PerCP	Peridinin chlorophyll

PMA	Phorbol 12-myristate 13-acetate
PPAR	Peroxisome proliferator activated receptor
PRR	Pathogen recognition receptor
QS	Quorum sensing
RANTES	Regulated upon activation, normally T cell expressed and secreted
RAR	Retinoic acid receptor
RXR	Retinoic X receptor
RNA	Ribonucleic acid
ROS	Reactive oxygen species
RPMI	Roswell Park Memorial Institute medium
RSG	Rosiglitazone: PPAR γ agonist
STAT	Signal transducer and activators of transcription
SD	Standard deviation
TGF- β	transforming growth factor beta
Th1	Type 1 helper T cell
Th2	Type 2 helper T cell
Th17	Type 17 helper T cell
TLR	Toll-like receptor
TNF	Tumor necrosis factor
Treg	Regulatory T cells
U	Units
V450	Violet 450 nm
Vs	Versus
YPD	Yeast peptone dextrose
α	Alpha
β	Beta
γ	Gamma
μ M	Micromolar

11 Index of tables and figures

Tables

Table 1. Antibodies used for cell surface molecules and intracellular staining analysis	25
--	----

Figures

Figure 1. TLR signaling pathway	9
Figure 2. Dectin-1 and dectin-2 signaling pathway	12
Figure 3. Nuclear receptor structure	13
Figure 4. Nuclear receptor mechanism of action	15
Figure 5. Generation of monocytes-derived DC.....	23
Figure 6. Viability of iDC differentiated in presence of FOH.....	28
Figure 7. TLR4 expression in FOH-differentiated iDC and mDC.....	30
Figure 8. CD40, CD80, CD86 and HLA-DR expression in FOH-differentiated DC activated with distinct stimuli.....	31
Figure 9. Cytokines and chemokines secretion of FOH-differentiated DC in response to distinct stimuli.....	32
Figure 10. CD1 expression of DC differentiated in presence of FOH.....	34
Figure 11. Expression of nuclear receptors and their target genes in DC-differentiated in presence of FOH.....	35
Figure 12. Surface molecules expression on DC differentiated in presence of farnesol and nuclear receptors agonists	37
Figure 13. Viability of DC differentiated with nuclear receptor antagonists.....	38
Figure 14. CD1 molecule expression induced by FOH in presence of nuclear receptor antagonists.....	40
Figure 15. CD1 molecule expression induced by FOH in presence of RXR α antagonist ..	41
Figure 16. CD80 and CD40 molecule expression modulated by FOH in presence of nuclear receptors antagonists	43
Figure 17. CD86 and HLA-DR molecule expression modulated by FOH in presence of nuclear receptors antagonists	44

Figure 18. Cytokines and chemokines release by DC differentiated in presence of FOH and NR antagonists.....	46
Figure 19. Surface molecule expression in FOH-differentiated mDC in presence of MAPK and NF- κ B inhibitors.....	48
Figure 20. Cytokine and chemokine release by DC differentiated in presence of FOH and NF- κ B and MAPK pathway inhibitors.....	50
Figure 21. Analysis of NF- κ B activation in FOH-differentiated mDC	51
Figure 22. Analysis of p38 activation in FOH-differentiated mDC	52
Figure 23. iNKT expansion by FOH treated-DC.....	54
Figure 24. IFN- γ production by T cells co-cultured with mDC	55
Figure 25. Diminished capacity to induce iNKT expansion by FOH-differentiated mDC is dependent on the altered secretion of IL-12 and IL-10	57
Figure 26. Altered release of IL-12 and IL-10 by FOH-treated mDC impacts their capacity to induce T helper 1 polarization	58
Figure 27. FOH enhances the surface expression of inhibitory receptors on mDC	60
Figure 28. FOH altered DC capacity to induce FOXP3+ regulatory T cells expansion.....	61
Figure 29. Farnesol modulates DC function through a coordinated activation of several signaling pathways.....	69

12 Appendix

12.1 Curriculum Vitae

Wolfgang Vivas

Date of birth: January 18, 1990
 Place of birth: Caracas, Venezuela
 Nationality: Venezuelan

Education

Dr. rer. nat. **Friedrich Schiller University Jena, Jena - Germany (2015-2020)**

Licenciado en Biología
(Diploma in Biology) **Universidad Central de Venezuela, Caracas - Venezuela (2007 - 2013)**

Research Experience

2015 –2019

Doctoral Researcher

Project: „Modulation of dendritic cell function by the fungal quorum sensing molecule farnesol”

Supervisor: Prof. Oliver Kurzai.

- Phenotypic and functional characterization of dendritic cells differentiated in presence of farnesol
- Determined the contribution of nuclear receptors in the expression of surface molecules and cytokines release of dendritic cells treated with farnesol
- Examined the role of MAPK and NF-κB signaling pathways in the altered functionality of dendritic cells stimulated with farnesol
- Evaluation of T cells activation by farnesol-instructed dendritic cells

2014 – 2015

PhD student

Project: “Role of protease-activated receptors in hemostasis”

Supervisor: Dr. Mercedes Lopez

- Isolation and culture of natural killer T cells from healthy donors
- Determination of cytokines from natural killer T cells stimulated with protease-activated receptors agonists

2010 – 2013:

Undergraduate thesis student

Project: “Role of regulatory T cells in the immune response against BCG in HIV-infected children under HAART treatment”

Supervisor: Prof. Miguel Alfonzo

- Isolation and culture of T cells and regulatory T cells from HIV-infected children under HAART treatment
- Determined the impact of HAART treatment in the phenotypic and functional properties of CD4⁺ and CD8⁺ T cells against BCG from HIV-infected children

Research Supports

2015 – 2018

PhD fellowship from the Jena School of Microbial Communication (JSMC)

2014 – 2015

PhD scholarship from the Venezuelan Institute for Scientific Research (IVIC)

Committee Leadership

2016 – 2018

JSMC doctoral researchers' representative

2011 – 2013

Biology student's representative at the faculty board

Awards

2014

Venezuelan representation and winning team of the Novartis International Biotechnology Leadership Camp 2014

2013

Thesis dissertation with honors. Universidad Central de Venezuela

Publications

- Wolfgang Vivas, Ines Leonhardt, Kerstin Hünninger, Antje Häder, Alessandra Marolda, Oliver Kurzai. (2019). *Multiple Signaling Pathways Involved in Human Dendritic Cell Maturation Are Affected by the Fungal Quorum-Sensing Molecule Farnesol*. **J Immunol**, 2019, 2013 (11) 2959-69.
- Alessandra Marolda, Kerstin Hünninger, Sarah Böttcher, Wolfgang Vivas, Jürgen Löffler, Marc Thilo Figge, Oliver Kurzai. *Candida species-dependent release of IL-12 by dendritic cells induces different levels of NK cell stimulation*. **J Infect Dis**. 2020, 2060-71.

Selected Presentations

Oral

- **52th Wissenschaftliche Tagung der Deutschsprachigen Mykologischen Gesellschaft e. V.** Innsbruck, Austria. 06 - 08.09.2018. *Candida albicans* quorum sensing molecule farnesol activates nuclear receptors and modulates dendritic cells functionality. Wolfgang Vivas, Ines Leonhardt, Oliver Kurzai
- **7th International Conference on Microbial Communication for Young Scientists.** Jena, Germany, 19 - 22.03.2018. *Candida albicans* quorum sensing molecule farnesol modulates human dendritic cells function through the PPAR γ pathway. Wolfgang Vivas, Ines Leonhardt, Oliver Kurzai

Poster

- **20th Congress of the International Society for Human and Animal Mycology.** Amsterdam, Netherlands. 30.06.2018 - 04.07.2018. *Candida albicans* quorum sensing molecule farnesol modulates human dendritic cells function by nuclear receptors activation. Wolfgang Vivas, Ines Leonhardt, Oliver Kurzai
- **47th Jahrestagung der Deutschen Gesellschaft für Immunologie.** Erlangen, Germany. 12 – 15.09.2017. The fungal quorum sensing molecule Farnesol modulates CD1 molecules expression in dendritic cells. Wolfgang Vivas, Ines Leonhardt, Oliver Kurzai
- **Annual Conference of the Association for General and Applied Microbiology (VAAM).** Jena, Germany. 13 – 16.03.2016. The fungal quorum sensing molecule farnesol impairs dendritic cells maturation. Wolfgang Vivas, Ines Leonhardt, Oliver Kurzai

Spoken Languages

Spanish (native proficiency)

English (C2 level/proficient user)

German (A2 level/basic user)

12.2 Acknowledgments

I want to thank everyone who made this possible. This is not only my achievement, but also yours.

First, I want to thank my supervisor Prof. Oliver Kurzai, for his support throughout my work in his group, for believing and trusting me with this project. Your decision of choosing me among other candidates has an unimaginable impact in my life, thanks to you I had the opportunity to meet a different culture and a great group of people. Not only you helped me professionally, but personally in times of need. You are a great inspiration for me and the definition of what a good person is.

To my lab friends. Big thanks to Dr. Ines Leonhardt and Dr. Kerstin Hünninger for your supervision during my lab work. Thank you for being always willing to help me with fruitful discussions, taking care of me and helping with my integration into the lab. Alessandra, for being an awesome colleague with a great sense of humor and delicious cooking, but mostly for being a support during our long hours in the lab. Sophia, for being a great friend in which I could always rely on, for your kindness and being always willing to help me with any problem I had. Antje, for being the most awesome colleague I could have asked for, thank you so much for the great times we had during our Buffy Coats isolation, and for always laughing of my bad jokes. Lysett, for being a great person and fostering a positive atmosphere in the office. Amelia, for your kindness in helping me during the last stages of my dissertation. Silke and Katrin, for helping me with everything related to administrative procedures and your kindness during my integration into the group and Germany.

I would like to extend my gratitude to the members of the Host Fungal Interfaces group, Slavena, Enrico, Bettina, Daniel, Phillip, Christin, and Franziska for being such a wonderful group of people who made my time in Septomics even better.

To Prof. Marc Thilo Figge for being available for any consultation during my studies and for helping me during the JSMC committees.

To my friends Naim, Angela, Jens, Prasad, Tilo, Sarah, Kathy, Andres, Sindy, Michael, Prerna, Martha, Jofran and Franz, for being always there, for taking care of me and sharing my burdens. I love you guys.

I would like to thank the Jena School for Microbial Communication, specially Katja and Hendrik for your kind assistance during administrative processes which guided me into life in Jena.

My parents in law for being the best ones anyone could have. Thank you very much for your kindness and making me a part of your family.

My brothers, for always being there for me and being an example of what good men are. My nephews for making me happy, I wish you a world full of happiness and opportunities.

I want to especially thank my parents because all your support, kindness and love made me the person who I am today. Even though it was not always easy, you kept believing in me and I will always be thankful to you. For me, you are an infinite source of inspiration.

Finally, my wife Dolly Montaña. Words cannot express my gratitude to you. Thank you for being my best friend, the one I can trust, laugh with, cry with, and lean on. Your unconditional support, words of encouraging, and love gave me the strength that I needed during this time. You make me happy and a better man, and I can only hope to live more beautiful moments by your side. I love you.

12.3 Declaration of originality

I herewith declare that I am familiar with the relevant course of examination for doctoral Candidates of the Faculty of Biological Sciences of the Friedrich Schiller University, Jena. I declare that I, personally, composed and wrote this dissertation and that I have acknowledged any and all additional assistance, personal communications and sources according to the rules of academic work. I declare that assistance provided by specific individuals during the study and writing of this dissertation has been indicated in full. I declare that I did not enlist any assistance of a doctoral consultant and that no third parties have received, either directly or indirectly, monetary benefits from me for work connected to this submitted dissertation. I declare that this dissertation has not been submitted as an examination paper for a state, or other, scientific examination. I also declare that I did not submit the same, a substantially similar or a different dissertation to another education body.

I am aware that a false declaration will have legal consequences.

.....

Wolfgang Vivas